arXiv:0909.3347v2 [hep-th] 8 Mar 2010

Correlation functions of disorder fields and parafermionic currents in Z_N Ising models

V. A. Fateev, Y. P. Pugai

L. D. Landau Institute for Theoretical Physics, RAS,
117940 Moscow, Russia
and
Laboratoire de Physique Théorique et Astroparticules,
CNRS-UM2, UMR 5207,
Université Montpellier II, Pl. E. Batallion,
34095 Montpellier, France

Dedicated to the memory of Aliosha Zamolodchikov.

Abstract

We study correlation functions of parafermionic currents and disorder fields in the Z_N symmetric conformal field theory perturbed by the first thermal operator. Following the ideas of Al. Zamolodchikov, we develop for the correlation functions the conformal perturbation theory at small scales and the form factors spectral decomposition at large ones. For all N there is an agreement between the data at the intermediate distances. We consider the problems arising in the description of the space of scaling fields in perturbed models, such as null vector relations, equations of motion and a consistent treatment of fields related by a resonance condition.

1 Introduction

The calculation of correlation functions is one of the most interesting problems in the two dimensional integrable quantum field theory. Complete exact solutions of this problem were found for free field models, as, for example, Ising model, and for the Conformal Field Theories (CFT).

In massive integrable field theories an elegant way of studying behaviors of two point correlation functions was proposed by Al. Zamolodchikov in Ref. [1]. In the case of the perturbed Lee-Yang model he studied short and long distance asymptotics of the two point correlation function of primary fields. The short distance behaviors of correlation function were investigated by developing the infra red safe perturbation theory [1], based on the knowledge of the exact Vacuum Expectation Values (VEVs) of local fields found by non-perturbative methods [2, 3, 4]. The correlators in the infra red region were given by using a form factor spectral decomposition [5, 6]. A very good agreement between the asymptotics at the intermediate distances $Mr \sim 1$ allowed to claim that correlation functions in Lee-Yang Model are effectively described in this approach at all distances. A further development has shown that the proposed in Ref. [1] method is a simple and effective tool for an analysis of basic properties of correlation functions in different integrable massive models.

In the present paper we continue our studying [7, 8] of scaling fields in the parafermionic CFT [9, 10] with the central charge

$$c = 2(N-1)/(N+2)\,, \quad N = 2, 3, \cdots\,, \tag{1}$$

perturbed by the first thermal operator ε_1

$$\mathcal{A} = \mathcal{A}_{CFT} + \lambda \int d^2x\; \varepsilon_1(x)\,. \tag{2}$$

The resulting theory is integrable and Z_N symmetric. Depending on the sign of λ the system is in the ordered or disordered phase. We fix $\lambda > 0$ phase where the $\tilde{Z}_N$ symmetry is destroyed and vacuum expectation values of disorder operators are non-zero. In this scaling model [11, 12], we study, within the approach of Ref. [1], the correlation functions of disorder operators and parafermionic currents. These objects, as well as correlators of some W-algebra descendants, are not so easy for a direct investigation, mainly because of the resonances, which appear in the construction. Namely, following the procedure of Ref. [1], we need to study the situations, where the scaling dimensions D_a and D_b of some fields $\mathcal{O}_a$ and $\mathcal{O}_b$ satisfy the condition

$$D_a = D_b + 2\,n\,(1 - \Delta_{\varepsilon_1}), \quad n > 0\,. \tag{3}$$

In this case, the field $\mathcal{O}_a$ has the n-th order resonance with the field $\mathcal{O}_b$ and there is an ambiguity in defining the renormalized field $\mathcal{O}_a$

$$\mathcal{O}_a \to \mathcal{O}_a + \mathrm{const}\,\lambda^n\,\mathcal{O}_b\,. \tag{4}$$

This typically results in a logarithmic scaling of the field $\mathcal{O}_a$ [4]. We observed that according to our general formulae the form factors of fields satisfying (3) formally coincide as functions of rapidities. This creates a problem of defining form factors of such scaling fields, since it is expected from the general settings [13, 14], that there is a one-to-one correspondence between conformal and scaling fields. We suppose to discuss a general prescription for form factors of the fields possessing resonances in a separate publication. In this paper we consider some examples of the phenomena.

From the other side, studying short distance behaviors of the correlators, we found, that the terms in the perturbative expansion, which correspond to fields with the condition (3), formally diverge for integer parameters N. To proceed further with these cases, we use the fact, that our exact expressions are defined for the models with arbitrary number N. We provide an analytic continuation over this parameter and obtain finite results for the correlators. We check, that our prescription leads to correct expressions for the Ising model ($N = 2$ case) [15], as well as for other known cases.

A matching between the long and short distance asymptotics, which we found for all N, can also be considered as an additional confirmation, supporting the consistency of the proposed form factors, the expressions for the short distance expansions, for VEVs and for the normalizations of the scaling fields.

We choose for studying the Z_N invariant Ising model, since it has several nice properties. It is related with a statistical system, which have a simple and clear description [9, 16] and many important physical applications, while the operators of this lattice model, in general, have an interesting quasi-locality property [17, 18] in the scaling limit. We recall the notions of the lattice Z_N model in the section 2. In the critical points the lattice model is described by the parafermionic CFT with the central charge (1). In section 3, we introduce necessary definitions and collect basic facts about this CFT, concentrating attention on the algebraic structure of the space of conformal fields. This information is essential for further analysis of the space of form factors of local fields in the case of massive integrable model, which we consider in the section 4 as the model of QFT, defined as scattering theory with the simple Z_N symmetric S matrix [12]. We discuss an algebraic prescription for the form factors of the scaling fields. In particular, we consider the actions of deformed parafermionic currents in the space of form factors and study related questions, such as deformation of the quantum equations of motion, null vector relations and a prescription for form factors of fields, satisfying the resonance condition (3). Finally, the correlation functions of scaling fields of the massive integrable model (2) are discussed in the section 5. We concentrate our attention to the conformal perturbation theory and demonstrate, how to apply the quantum equation of motion to the computing coefficients of the perturbation theory. We discuss the regularization prescription for the resonances, appearing in the perturbation theory and also provide results of numerical computations.

2 Lattice Z_N-Ising model

In this section we recall following Ref. [9] basic definitions of the two dimensional lattice model, generalizing the well-known Z_2-symmetric Ising model to the case with the Z_N, $\{N = 2, 3, \ldots\}$ symmetry.

We consider the model of the statistical mechanics defined at the square lattice. Let spin variables σ are associated with the cites of the lattice and take values in the group Z_N, i.e. $\sigma = \omega^k$ $(k = 0, \ldots, N-1)$, where $\omega = \exp\left(\frac{2\pi i}{N}\right)$.



Fig 1. Z_N Ising model at the lattice.

We study the local theory where Boltzmann weight $e^{-\mathcal{H}(\sigma,\sigma')} = \mathcal{W}(\sigma,\sigma')$ depends on spins σ and σ' situated at neighboring sites. The partition sum of the model is, by definition

$$\mathcal{Z} = \sum_{spins} \prod_{edges} \mathcal{W}(\sigma, \sigma')\,. \tag{5}$$

In Z_N Ising model Boltzmann weights satisfy the Z_N symmetry $\mathcal{W}(\sigma, \sigma') = \mathcal{W}(\omega\sigma, \omega\sigma')$, and the reality condition $\mathcal{W}(\sigma, \sigma') = \mathcal{W}(\sigma^\dagger, \sigma'^{\,\dagger})$. It means, that, up to a normalization constant, the function $\mathcal{W}(\sigma, \sigma')$ has the form

$$\mathcal{W}(\sigma, \sigma') = \sum_{k=0}^{N-1} \mathcal{W}_k (\sigma^\dagger \sigma')^k, \qquad \mathcal{W}_0 = 1\,, \tag{6}$$

where the parameters $\mathcal{W}_k$ of the model are real non-negative numbers satisfying the equation $\mathcal{W}_k = \mathcal{W}_{N-k}$.

An important information about the model is encoded in a set of its correlation functions. For example, the correlation functions of the spin operators are defined at the lattice as

$$\langle \sigma_{k_1}(x_1) \cdots \sigma_{k_s}(x_s) \rangle = \frac{1}{\mathcal{Z}} \sum_{spins} \sigma^{k_1}(x_1) \cdots \sigma^{k_s}(x_s) \prod_{edges} \mathcal{W}(\sigma, \sigma')\,.$$

We will study a system in the thermodynamic limit, assuming appropriate periodic conditions at the infinity. The one point correlation functions $\langle \sigma_k \rangle$ serve as a measure of order at the system. Another important lattice operators are the disorder operators [10, 17, 18]. Consider a directed path $\mathbf{\Gamma}$ going through points of a dual lattice and intersecting the bonds of the original lattice. Let all weights $\mathcal{W}_s$ at the bonds, crossed by path $\mathbf{\Gamma}$, are changed to become $\tilde{\mathcal{W}}_s^{(k)} = \mathcal{W}_s\ \omega^{sk}$. The presence of the dislocation along the path introduces in the system a fractional domain wall, that favors a discontinuity in the value of the neighboring spins by k. The partition function $\tilde{\mathcal{Z}}_{\mathbf{\Gamma}}$ on the inhomogeneous lattice now differs from the original one. We interpret the dislocation as an insertion of two operators μ_k and $\mu_k^\dagger = \mu_{N-k}$ situated at the sites of the dual lattice as it is shown in Fig.2.



Fig 2. Two point correlator of disorder operators μ_k and $\mu_k^\dagger$.

By definition, the two point correlation function of conjugate disorder operators is given as

$$\langle \mu_k(x_1) \mu_k^\dagger(x_2) \rangle_{\mathbf{\Gamma}} = \frac{1}{\mathcal{Z}}\ \tilde{\mathcal{Z}}_{\mathbf{\Gamma}}\,.$$

Here $x_{1,2}$ are the coordinates of the corresponding operators. This interpretation turns out to be useful since the dependence on the contour $\mathbf{\Gamma}$ is not very strong. As in the Ising model [17], one can make contour deformations by closed paths, without changing the correlation function. Applying this freedom, we use prescription of attaching contours from infinity to each of the points x_j of the dual lattice, as it is shown in Fig.3. Now, the definition of two point correlation functions can be immediately extended to the multi-point case, including disorder operators with a total non-zero Z_N charge.



Fig 3. Correlation function $\langle \mu_k \ \mu_{N-k} \rangle$.

In a general case of correlators of spin and disorder operators, only an absolute value of the correlation function remains to be a path independent, and one should fix relative position of contours [10]. This is because, the correlation functions of the product of spin and disorder operators, before and after a complete counterclockwise rotation of the disorder variable around order variable, differ by a phase ω^{-kl}. Equivalently, this happens, whenever a μ_k-path crosses a σ_l variable. Operators with that properties are called mutually quasi-local with the exponent $\gamma_{kl} = -kl/N$.

Order and disorder parameters are basic operators in the theory. Other operators are constructed as their operator products. In general, if the distance $|x - y|$ between operators is much smaller than the correlation length, then the local operators $\psi_{l,k}$, appearing at the operator product

$$\sigma_l(x)\mu_k(y) = C_{lk}(x-y)\psi_{l,k}(x) + \cdots ,$$

do obey the parastatistics [18]. In particular, further we will study the correlation functions of two parafermionic currents $\psi = \psi_{1,1}$ and $\psi^\dagger$ in the scaling limit.

Under the Krammers Wannier duality [17] order parameters become disorder parameters and wiseverse. The hyperplane of self-duality has the dimension $\left[\frac{N}{4}\right]$. For the Ising model ($N = 2$ case) and three state Potts model ($N = 3$ case), the system has a second order phase transition at the self-dual point. For $N = 4$ it coincides with the well-known Ashkin-Teller model. For prime $N \geq 5$, the Z_N theory has, as a rule, three phases (ordered, disordered and Kosterlitz-Thouless phases). For non-prime N the phase structure is more complicated. As an example, we consider the phase diagram of the Z_5 model [16].



Fig 4. The phase diagram of the Z_5 model

Here the phase I, the ordered phase, is characterized by the conditions $\langle\sigma_l\rangle \neq 0$ and $\langle\mu_k\rangle = 0$. In the disordered phase II the situation is inverse $\langle\sigma_l\rangle \neq 0$ and $\langle\mu_k\rangle = 0$. Finally, in the Kosterlitz-Thouless phase III expectation values of operators of both type are zero $\langle\sigma_l\rangle = \langle\mu_k\rangle = 0$. The line FB denotes the self-duality region. It contains two symmetrically situated "bifurcation points" **C** and **C***. Along the line **CC*** the model have a first order phase transition and ordered and disordered phases may coexist. The points **C** and **C*** are integrable and critical [28, 16]. The theory in these points has $Z_N \times \tilde{Z}_N$ symmetry and is described by the continuous parafermionic CFT constructed in Ref. [9]. In the scaling limit in the vicinity of a critical point the order σ_l and disorder μ_k operators, as well as parafermions ψ, are described by the fields, depending on the continuous space parameters. We preserve for that fields the same notations as on the lattice.

Due to the conformal invariance and infinite-dimensional symmetry of the critical theory [19], the analysis of its correlation functions, as well as the structure of its space of states, simplify drastically. This was described in details in Refs. [9, 10]. Basing on CFT results, one can study basic properties of the correlation functions of the Z_N models in a vicinity of the critical point by application of the conformal perturbation theory. Again, the easiest for an analysis perturbations are those, which are integrable, due to a presence of an infinite set of integrals of motion [13]. Different integrable perturbations of the conformal field theories [9] were studied in Ref. [11].

In this paper, we study a vicinity of the critical point **C** in the phase II, where the temperature deformation leads to an appearance of the finite correlation length and to non-zero vacuum expectation value of the disorder parameters [11, 12, 20]. In the QFT language, that is the massive perturbation of parafermionic CFT (2) by the most relevant first thermal operator ε_1, which destroys the dual $\tilde{Z}_N$ symmetry and preserves the symmetry Z_N.

One of the questions, which we would like to address in our study, is a structure of the space of scaling fields. We expect, that the space of composite quasi-local fields constructed from operators σ_k and μ_l in the vicinity of critical point, will be essentially the same, as in the CFT. Let us note, that this statement was supported by several results on counting of local operators in various integrable models [14, 22]. To understand this problem, we tried to apply in Ref. [7, 8] a knowledge on the algebraic structure of lattice operators, which is based on a deformation of conformal algebras [23, 24, 25, 26]. According to this idea we recall first the structure of quasi-local fields in the CFT point. Then, we will try to apply the clear and simple algebraic scheme for the investigations of matrix elements of scaling fields in the basis of asymptotic states and also for studying their correlation functions off-criticality.

3 Space of states in parafermionic CFT

In this section, we recall basic facts [9, 10] about the conformal field theory with the parafermionic symmetry which describes critical points of Z_N Ising model [9] and related models [32].

3.1 Parafermionic symmetry

In the conformal limit the order σ_k and disorder μ_k parameters, which determine long range correlations of spins and dual spins, have the anomalous dimensions

$$2d_k = \frac{k(N-k)}{N(N+2)}\,. \tag{7}$$

Their Z_N and dual $\tilde{Z}_N$ charges, respectively, are equal to k. Under the action of the Z_N symmetry spin fields transform as

$$\sigma_k \to \omega^{kn}\sigma_k, \quad n \in Z\,. \tag{8}$$

The transformation law for operators μ_k under the action of the dual $\tilde{Z}_N$ symmetry looks similarly

$$\mu_l \to \omega^{ln'}\mu_l, \quad n' \in Z\,. \tag{9}$$

The spin and disorder fields are the basic operators in the theory. All other fields are constructed from them. The composite fields are naturally separated into families with the fixed additive $Z_N \times \widetilde{Z}_N$ charges (k, l). The members of the family behave under $Z_N \times \widetilde{Z}_N$ transformation as

$$\Phi \to \omega^{kn+ln'}\Phi, \quad n, n' \in Z\,.$$

The parafermionic fields ψ_k and $\overline{\psi}_k$ generalizing the usual Ising model fermions appear in the OPE of the order and disorder fields

$$\begin{aligned}\sigma_k(z,\overline{z})\mu_k(0,0) &= |z|^{-4d_k} z^{\Delta_k}[\ \psi_k(0) + \cdots]\,,\\ \sigma_k(z,\overline{z})\mu_k^{\dagger}(0,0) &= |z|^{-4d_k} \overline{z}^{\Delta_k}[\ \overline{\psi}_k(0) + \cdots]\,.\end{aligned} \tag{10}$$

These currents are holomorphic and generate the infinitely dimensional symmetry due to the conservation laws

$$\partial_{\overline{z}}\psi_k = 0\,, \qquad \partial_z\bar{\psi}_k = 0\,. \tag{11}$$

We concentrate our attention to the simplest solution of the associativity condition for the operator algebra of currents, which corresponds to CFT with the central charge (1), and the conformal dimensions of currents

$$\Delta_k = \frac{k(N-k)}{N}\,. \tag{12}$$

The fields $\psi = \psi_1$ (and respective antichiral currents) are the basic ones in the parafermionic algebra. It is convenient for us to consider, as well, conjugate currents $\psi_{N-1} = \psi^\dagger$.

In the conformal model the space of states splits naturally into a direct sum of subspaces with the specified $Z_N \times \widetilde{Z}_N$ charge (k,l)

$$\{F\} = \oplus\, \{F\}_{[m,\overline{m}]}, \quad N \geq m, \overline{m} \geq 1-N\,, \tag{13}$$

where $[m,\overline{m}] = [k+l, k-l]\,, m+\overline{m} \in 2Z$. In these notations parafermionic currents and order-disorder fields belong to the following subspaces

$$\begin{aligned} \psi \in \{F\}_{[2,0]}\,, \quad & \psi^\dagger \in \{F\}_{[-2,0]}\,, \\ \overline{\psi} \in \{F\}_{[0,2]}\,, \quad & \overline{\psi}^\dagger \in \{F\}_{[0,-2]}\,, \\ \sigma_k \in \{F\}_{[k,k]}\,, \quad & \mu_k \in \{F\}_{[k,-k]}\,. \end{aligned} \tag{14}$$

Conformal fields are classified according to the representations of the parafermionic algebra. The action of the parafermionic generators A_ν $(A^\dagger_\nu)$ is defined by the OPE

$$\begin{aligned} \psi(z)\Phi_{[m,\bar{m}]} &= \sum z^{-\frac{m}{N}+n-1} A_{\frac{1+m}{N}-n}\; \Phi_{[m,\bar{m}]}\,, \\ \psi^\dagger(z)\Phi_{[m,\bar{m}]} &= \sum z^{\frac{m}{N}+n-1} A^\dagger_{\frac{1-m}{N}-n}\; \Phi_{[m,\bar{m}]}\,. \end{aligned} \tag{15}$$

Notice, that, if $\Phi_{[m,\bar{m}]} \in F_{[m,\bar{m}]}$ has the conformal dimensions $(d,\bar{d})$ then the conformal dimensions of fields

$$A_\nu \Phi_{[m,\bar{m}]} \in \{F\}_{[m+2,\bar{m}]}\,, \quad A^\dagger_\nu \Phi_{[m,\bar{m}]} \in \{F\}_{[m-2,\bar{m}]}\,,$$

are $(d-\nu, \bar{d})$.

Order and disorder fields are the primaries of the parafermionic algebra. For instance, the following equations hold for $n \geq 0$

$$\begin{aligned} A_{\frac{1+k}{N}+n}\; \mu_k &= A^\dagger_{\frac{1-k}{N}+n+1}\; \mu_k = 0\,, \\ \bar{A}_{\frac{1-k}{N}+n+1}\; \mu_k &= \bar{A}^\dagger_{\frac{1+k}{N}+n}\; \mu_k = 0\,. \end{aligned} \tag{16}$$

All other fields of the model are obtained by the action of the currents $\psi, \bar{\psi}$ on the fields μ_k. The space of states of the CFT decomposes into a direct sum of irreducible representations of the parafermionic algebra

$$\{F\} = \oplus_{k=0}^{N-1}\; [\mu_k]_{A,\bar{A}}\,. \tag{17}$$

This space can be also obtained by an application of parafermionic generators to the order parameters σ_k. Moreover, in some situations we will need another infinite symmetry description of parafermionic CFT, namely, the W symmetry, which we recall in the next subsection.

3.2 W algebra symmetry

The space of fields in parafermionic CFT allows a classification with respect to another infinite dimensional symmetry algebra, the so-called W_N algebra [27]. The generators of the latter $W_2(z),\ W_3(z), \ldots$ appear at the operator product of parafermionic currents

$$\psi(z)\psi^\dagger(0) = \frac{1}{z^{2\Delta_1}}\Big(1 + z^2\frac{N+2}{N}W_2(0) + \\ + z^3\big(\frac{1}{N^{\frac{3}{2}}}W_3(0) + \frac{N+2}{2N}\partial W_2(0)\big) + \cdots\Big)\,. \tag{18}$$

Here W_2 currents with spin 2 generate Virasoro algebra with the central charge given by eq. (1). The currents W_3 have spin 3. They generate the whole algebra, including the higher spins currents, which are omitted in eq. (18) .

Currents of W algebra, and respective anti chiral currents, have zeroth Z_N charges. Acting on the highest weight fields, they create an irreducible representation. From the viewpoint of the $W_N \times \bar{W}_N$ symmetry, each of the spaces $[\mu_k]_{A,\bar{A}}$ expands into a direct sum of representations. Namely, let us denote as $(\psi^\dagger)^l\mu_k$ the field with the minimal conformal dimension, that can be obtained by the $l-$times application of the parafermionic generators $\psi^\dagger$ to μ_k

$$(\psi^\dagger)^l\mu_k = A^\dagger_{\frac{2l-1-k}{N}}A^\dagger_{\frac{2l-3-k}{N}}\cdots A^\dagger_{\frac{1-k}{N}}\mu_k\,. \tag{19}$$

Its conformal dimensions are easily computed from (7). Then, up to a normalization, the following relations take place

$$\begin{aligned} \Phi^{(k)}_{k-2l,-k+2\bar{l}} &= (\psi^\dagger)^l(\overline{\psi})^{\bar{l}}\mu_k\,, \quad l,\bar{l} = 0,1,...,k\,, \\ \Phi^{(k)}_{k+2l,-k-2\bar{l}} &= (\psi)^l(\overline{\psi}^\dagger)^{\bar{l}}\mu_k\,, \quad l,\bar{l} = 0,1,...,N-k\,. \end{aligned} \tag{20}$$

These fields $\Phi^{(k)}_{m,\bar{m}}$ are the W algebra primaries. The action of generators of W algebra on it create the irreducible representation $[\Phi^{(k)}_{m,\bar{m}}]_{W,\bar{W}}$. The explicit values of the conformal dimensions $(d^{(k)}_m, \bar{d}^{(k)}_m)$ of these fields are given as follows

$$d^{(k)}_m = \frac{(k+1)^2-1}{4(N+2)} - \frac{m^2}{4N}\,, \quad -m \le k \le m\,. \tag{21}$$

For the cases, when $|m| > k$, we use the relation $d^{(k)}_m = d^{(N-k)}_{m-N}$, which follows from the Z_N symmetry condition

$$\Phi^{(k)}_{m\bar{m}} = \Phi^{(N-k)}_{m-N,-N+\bar{m}}\,. \tag{22}$$

For example, the physically important energy fields $\varepsilon_k = \Phi^{(2k)}_{0,0}$ are among these primaries. Operators ε_k are local with respect to all fields and have the conformal dimensions

$$D_k = k(k+1)/(N+2). \tag{23}$$

We refer to the paper [9] for further details. Let us only comment here, that the Z_N symmetric parafermionic CFT can be equivalently considered as a particular case of the conformal field theory $\mathcal{WM}_N^p$, introduced in Ref. [27]. Namely, for the fixed value of N it has the smallest central charge (1) among all rational unitary minimal conformal theories with the extended W_N algebra, which corresponds to the parameter $p = N + 1$.

We will use the conformal data, represented above in the next sections, to describe the perturbed conformal operators, for which we again will preserve the same CFT notations.

4 Form factor approach

Now we turn to the operators in the corresponding massive integrable theory (2), which allows several equivalent descriptions. In the given section, we describe it as a two dimensional QFT model with the factorized scattering of Z_N charged particles. The particles $a \in \{1, \ldots, N-1\}$ in the Z_N $(N = 2, 3, 4, \ldots)$ symmetric models have masses [20]

$$M_a = M \frac{\sin(\pi a/N)}{\sin(\pi/N)} \,. \tag{24}$$

The antiparticle $a^\dagger$ is, by definition, identified with the particle $N - a$. The scattering matrix of the lightest particles $a = 1$ has a simple form [12]

$$S_{11}(\beta) = \frac{\sinh(\frac{\beta}{2} + \frac{i\pi}{N})}{\sinh(\frac{\beta}{2} - \frac{i\pi}{N})} \,. \tag{25}$$

The S matrices for higher particles are also diagonal and can be extracted from S_{11}, according to a standard bootstrap prescription. For example, the scattering matrix between the particle 1 and the antiparticle $1^\dagger$ is $S_{11^\dagger}(\beta) = S_{11}(i\pi - \beta)$.

The knowledge of the exact spectrum (24) and the scattering matrix (25) allows to study correlation functions of the theory, by using its spectral decomposition into the series of form factors. The form factors

$$\langle \mathcal{O}(x) | \beta_1, \ldots, \beta_n \rangle_{a_1, \ldots, a_n} \,, \tag{26}$$

of the scaling field $\mathcal{O}(x)$ are matrix elements of this operator in a basis of asymptotic states, formed by particle creation operators. We assume, that the particles, labeled by $a_1, \cdots, a_n$, have the rapidities $\beta_1, \cdots, \beta_n$. The functions (26) should satisfy to some analytical properties and, also, to a set of functional equations, the so-called form factor axioms [5, 6], to guarantee the (quasi)locality of scaling fields.

An usual problem in the form factor approach (see, for example [29, 30, 31] for Z_N Ising models case) is that it is not so easy to determine, which scaling field is described by the given functions, satisfying proper functional equations and analyticity conditions. Moreover, for a matching with short distance formulae, it is necessary to determine the normalization of the scaling fields, described by form factors. In our construction [7, 8], we proposed some algebraic approach to solve these problems. Still, there are many subtle questions in this direction. We would like to discuss some of them in the next sections.

4.1 Free fields construction for form factors

In Refs. [7, 8] we have followed the algebraic approach [33, 25, 23, 26, 34] to the form factors of scaling limit of the ABF model [32]. This lattice model falls into the same universality class, as the Z_N Ising model. In the corner transfer matrix approach its hidden symmetry is a deformation of CFT symmetry algebras (7)-(23). We explore algebraic maps in the space of form factors to produce exact expressions for form factors of scaling fields. Our basic prescription, derived in Ref. [7, 8] for form factors, can be shortly re-formulated as following. We introduce the notations [1]

$$\begin{aligned} \mathcal{B}^\dagger(\beta)^{(k)}_{m,\bar m} &= \frac{e^{\frac{m-\bar m}{2N}\beta}}{\sqrt{2\mathrm{sin}\frac{\pi}{N}}}\sum_{a=\pm} a\ e^{\frac{i\pi}{2N}(k+1-\frac{m-\bar m}{2})a}\mathcal{Z}^\dagger_a(\beta)\,, \\ \mathcal{B}(\beta')^{(k)}_{m,\bar m} &= -\frac{e^{-\frac{m-\bar m}{2N}\beta'}}{\sqrt{2\mathrm{sin}\frac{\pi}{N}}}\sum_{b=\pm} b\ e^{\frac{i\pi}{2N}(-k-1-\frac{m-\bar m}{2})b}\mathcal{Z}_b(\beta')\,. \end{aligned} \tag{27}$$

The explicit expressions for the form factors of fields $(\psi^\dagger)^l(\bar\psi)^{\bar l}\mu_k$ in the perturbed theory are given for $m = k-2l,\ \bar m = k-2\bar l$ as

$$\langle(\psi^\dagger)^l(\bar\psi)^{\bar l}\mu_k|\{\beta\},\{\beta'\}\rangle_{(n,n')} = C^{(k)}_{m,\bar m}\langle\!\langle\prod_1^n\mathcal{B}(\beta_j)^{(k)}_{m,\bar m}\prod_1^{n'}\mathcal{B}^\dagger(\beta'_j)^{(k)}_{m\bar m}\rangle\!\rangle\,. \tag{28}$$

We assume in this equation, that the form factor is zero, unless the Z_N neutrality condition $2(n'-n) = m+\bar m$ is satisfied. The label n stands for the number of lightest antiparticles $1^\dagger$, carrying the Z_N charge -2, and n' means the number of particles 1 with the Z_N charge 2. The constant $C^{(k)}_{m,\bar m}$ is determined by the normalization of the scaling field. In what follows we will specify its value for spin-less fields and for parafermionic currents to be in agreement with the conformal normalization. In eq. (28), we used a shorthand notation for the state with these numbers of particles and antiparticles

$$|\{\beta\},\{\beta'\}\rangle_{(n,n')} \equiv |\beta_1,\ldots,\beta_n,\beta'_1,\ldots,\beta'_{n'}\rangle_{1^\dagger\cdots1^\dagger,1\cdots1}\,. \tag{29}$$

The symbol of ordered product of particle creation operators in eq. (28) is used for the object

$$\prod^n\mathcal{B}(\beta_j) = \mathcal{B}(\beta_1)\cdots\mathcal{B}(\beta_n)\,, \tag{30}$$

---

[1] Matrix elements of operators $\mathcal{Z}_a,\ \mathcal{Z}^\dagger_a$ are defined by the rules (31)-(32) below. A more explicit definition of these operators can be found in Ref. [23].

which is a linear combination of the products of exponential free bosonic fields $\mathcal{Z}_{\pm}(\beta)$, see Ref. [23] for details. The expectation value of a product of operators $\mathcal{Z}_a(\beta)$ and $\mathcal{Z}_a^{\dagger}(\beta)$ over the Fock vacuum can be computed by applying the Wick theorem

$$\begin{aligned}
&\langle\!\langle \mathcal{Z}_{a_1}(\beta_1)\cdots\mathcal{Z}_{a_n}(\beta_n)\mathcal{Z}^{\dagger}_{b_1}(\beta'_1)\cdots\mathcal{Z}^{\dagger}_{b_m}(\beta'_m)\rangle\!\rangle \\
&=\prod_{i<j}\langle\!\langle \mathcal{Z}_{a_i}(\beta_i)\mathcal{Z}_{a_j}(\beta_j)\rangle\!\rangle\prod_{i<j}\langle\!\langle \mathcal{Z}^{\dagger}_{b_i}(\beta'_i)\mathcal{Z}^{\dagger}_{b_j}(\beta'_j)\rangle\!\rangle\times \\
&\quad\prod_{i,j}\langle\!\langle \mathcal{Z}_{a_j}(\beta_j)\mathcal{Z}^{\dagger}_{b_i}(\beta'_i)\rangle\!\rangle\,,
\end{aligned}\tag{31}$$

The contraction rules of two operators in this equation are determined in terms of the meromorphic functions $\zeta(\beta)$, $\zeta^{\dagger}(\beta)$, given in the appendix B, as following (we assume, that $\beta=\beta_1-\beta_2$)

$$\begin{aligned}
&\langle\!\langle \mathcal{Z}_a(\beta_1)\mathcal{Z}_b(\beta_2)\rangle\!\rangle=\langle\!\langle \mathcal{Z}^{\dagger}_{-a}(\beta_1)\mathcal{Z}^{\dagger}_{-b}(\beta_2)\rangle\!\rangle \\
&\qquad=\zeta(\beta)\frac{\sinh\bigl(\frac{\beta}{2}+\frac{i\pi}{2N}(a-b)\bigr)}{\sinh\frac{\beta}{2}}\,, \\
&\langle\!\langle \mathcal{Z}_a(\beta_1)\mathcal{Z}^{\dagger}_b(\beta_2)\rangle\!\rangle=\langle\!\langle \mathcal{Z}^{\dagger}_{-a}(\beta_1)\mathcal{Z}_{-b}(\beta_2)\rangle\!\rangle \\
&\qquad=\zeta^{\dagger}(\beta)\cosh\Bigl(\frac{\beta}{2}-\frac{i\pi}{2N}(a+b)\Bigr)\,.
\end{aligned}\tag{32}$$

4.2 Parafermionic currents action

Let us shortly comment the algebraic structures, encoded in the form factors prescription described above. We obtained the equation (28) starting from the thermal operators operator form factors case $m=-\bar{m}$ and $n'=n$ [23] as a result of the parafermionic currents action [7] on the bosonic operators

$$\begin{aligned}
&\mathcal{Z}^{\dagger}_a(\beta)\xrightarrow{(\psi)}\mathcal{Z}^{\dagger}_a(\beta)e^{\frac{\beta}{N}-\frac{i\pi}{2N}a}\,, &&\mathcal{Z}_b(\beta)\xrightarrow{(\psi)}\mathcal{Z}_b(\beta)e^{-\frac{\beta}{N}-\frac{i\pi}{2N}b}\,, \\
&\mathcal{Z}^{\dagger}_a(\beta)\xrightarrow{(\bar{\psi})}\mathcal{Z}^{\dagger}_a(\beta)e^{-\frac{\beta}{N}+\frac{i\pi}{2N}a}\,, &&\mathcal{Z}_b(\beta)\xrightarrow{(\bar{\psi})}\mathcal{Z}_b(\beta)e^{\frac{\beta}{N}+\frac{i\pi}{2N}b}\,, \\
&\mathcal{Z}^{\dagger}_a(\beta)\xrightarrow{(\psi)^{\dagger}}\mathcal{Z}^{\dagger}_a(\beta)e^{-\frac{\beta}{N}+\frac{i\pi}{2N}a}\,, &&\mathcal{Z}_b(\beta)\xrightarrow{(\psi)^{\dagger}}\mathcal{Z}_b(\beta)e^{\frac{\beta}{N}+\frac{i\pi}{2N}b}\,, \\
&\mathcal{Z}^{\dagger}_a(\beta)\xrightarrow{(\bar{\psi})^{\dagger}}\mathcal{Z}^{\dagger}_a(\beta)e^{\frac{\beta}{N}-\frac{i\pi}{2N}a}\,, &&\mathcal{Z}_b(\beta)\xrightarrow{(\bar{\psi})^{\dagger}}\mathcal{Z}_b(\beta)e^{-\frac{\beta}{N}-\frac{i\pi}{2N}b}\,.
\end{aligned}\tag{33}$$

Such prescription determines a set of maps in the space of multi-particle form factors, for example,[2]

$$\langle 0|\Phi^{(k)}_{m,\bar{m}}|\{\beta\},\{\beta'\}\rangle_{(n,n')}\xrightarrow{(\psi)}\langle 0|(\psi)\Phi^{(k)}_{m,\bar{m}}|\{\beta\},\{\beta'\}\rangle_{(n-1,n')}\,.\tag{34}$$

This relation, and similar relations for other parafermionic current actions, are understood as following. We assume, that the multi-particle form factors for the field $\Phi^{(k)}_{m,\bar{m}}$ are given as matrix elements of a linear combination of products of n operators $\mathcal{Z}_b(\beta)$ and n' operators $\mathcal{Z}^\dagger_a(\beta)$. Then, the form factors of the field $(\psi)\Phi^{(k)}_{m,\bar{m}}$ will be given by the modified bosonization prescription, where the number of antiparticles is one less, and the change (33) is provided for each of $\mathcal{Z}_a$ and $\mathcal{Z}^\dagger_a$ operators. One can see, that this prescription obviously agree with the equation (28). The eq. (33) was obtained from the deformed parafermionic action. For this reason, the following identification was proposed for $l, \bar{l} = 0, 1, ..., k$ in Ref. [7]

---

[2] We use in eq. (34) the conformal notations $(\psi)\Phi^{(k)}_{m,\bar{m}}$, since this equation is a perturbed analogue of eq. (19).

$$\langle 0|\Phi^{(k)}_{k-2l,-k+2\bar{l}}|\{\beta\},\{\beta'\}\rangle_{(n,n')} = \langle 0|(\psi^\dagger)^l(\overline{\psi})^{\bar{l}}\mu_k|\{\beta\},\{\beta'\}\rangle_{(n,n')} \,. \quad (35)$$

The validity of this equation for multi-particle form factors with $\bar{m} = -m$ was checked by comparing a prescription for the form factors (27)-(33) with the form factors of the scaling fields in the left hand side of this equation, obtained in the prescription for the deformed W algebra primaries [33]. We also get a clear evidence on the correctness of this identification by studying the correlation functions of the order and disorder fields in Ref. [7].[3] We consider the equation (35) as an off critical analogue of the equation (20).

It is possible to continue further a studying of the structure of the space of form factors of scaling fields in an analogy with the correspondent algebraic description of CFT. We note here, that, together with the equation (20), the form factor prescription (27)-(33) also satisfy the charge conjugation condition (22). Using the Z_N symmetry condition $\psi = \Phi^{(N)}_{N-2,-N}$, it is possible to derive the form factors of the parafermionic currents. A simple observation is that these matrix elements are related with the form factors of the field $\Phi^{(2)}_{2,0} = (\psi)\varepsilon_1$ as following (see also [29])

$$\langle\psi|\{\beta\},\{\beta'\}\rangle_{(n-1,n)} = \lambda\sqrt{\frac{2}{N}}\Big(\sum^{n-1} e^{-\beta_j} + \sum^{n} e^{-\beta'_j}\Big)^{-1}\langle\Phi^{(2)}_{2,0}|\{\beta\},\{\beta'\}\rangle_{(n-1,n)} \,. \quad (36)$$

This equation is an off critical analogue of the quantum equation of motion (11) for the perturbed parafermionic currents

$$\frac{\partial}{\partial\bar{z}}\psi(z,\bar{z}) = \lambda\sqrt{\frac{2}{N}}(\psi)\epsilon_1(z,\bar{z}) = \lambda\sqrt{\frac{2}{N}}\Phi^{(2)}_{2,0}(z,\bar{z}) \,. \quad (37)$$

In the general multi particle case this equation follows from the trigonometric functions identities. Further we will use eq. (37) in the short distance expansion.

Our next comment on a mapping between form factors and interpreting our form factors as matrix elements of the scaling fields is as following. Formally applying the prescription of Ref. [7], we arrive to fact, that the null vectors equation (16) survives after the perturbation (see Ref. [22], where a similar phenomena was studied in the sine-Gordon model context). It can be directly checked, that for perturbed fields we have relations

---

[3] Moreover, we demonstrated in Ref. [7], that the deformed parafermionic currents, acting on the deformed vertex operators, reproduce also VEVs for perturbed W algebra primaries, and, in particular, explain the factorized form of resulting VEVs (56) of the fields $\Phi^{(k)}_{m,\bar{m}}$.

$$A_{\frac{1+k}{N}}\ \mu_k = 0\,, \quad \bar{A}^{\dagger}_{\frac{1+k}{N}}\ \mu_k = 0\,. \tag{38}$$

The first equation here follows from the parafermionic currents action procedure, introduced in Ref. [7]. Indeed, one can check, that taking the $\alpha \to 0$ limit in the expression

$$\begin{aligned}&\langle A_{\frac{1+k}{N}}\Phi^{(k)}_{k,-k}|\{\beta\},\{\beta'\}\rangle_{(n-1,n)} \sim \\ &\lim_{\alpha\to\infty} e^{-\frac{1+k}{N}\alpha}\langle\!\langle \prod^{n-1} \mathcal{B}(\beta_j)^{(k)}_{k,-k} \prod^{n} \mathcal{B}^{\dagger}(\beta'_j)^{(k)}_{k,-k}\ \mathcal{B}(\alpha)^{(k)}_{k,-k}\rangle\!\rangle\,,\end{aligned} \tag{39}$$

we get exact zero for arbitrary particle number n. For the second equation, after the application of the Wick theorem, we find that the condition

$$\langle 0|\bar{A}^{\dagger}_{\frac{1+k}{N}}\ \mu_k|\beta_1,\ldots,\beta_{s+1},\beta'_1,\ldots,\beta'_s\rangle = 0\,, \tag{40}$$

for the arbitrary $2s+1$ particle matrix element of the scaling field $\bar{A}^{\dagger}_{\frac{1+k}{N}}\mu_k$ is reduced to proving the following trigonometric identity

$$\begin{aligned}&\sum_{\{a_j\},\{b_j\}} \prod_{i=1}^{s+1} a_i \prod_{j=1}^{s} b_j\ e^{-\frac{i\pi}{N}(k+1)b_j} \prod_{i=1}^{s+1}\prod_{j=1}^{s} \cosh\Big(\frac{\beta_i-\beta'_j}{2} - \frac{i\pi}{2N}(a_i+b_j)\Big) \times \\ &\times \prod_{i<j}^{s+1} \sinh\Big(\frac{\beta_i-\beta_j}{2} + \frac{i\pi}{2N}(a_i-a_j)\Big) \prod_{i<j}^{s} \sinh\Big(\frac{\beta'_i-\beta'_j}{2} - \frac{i\pi}{2N}(b_i-b_j)\Big) = 0\,,\end{aligned}$$

which follows from equation (28).

Still, the null vector conditions, as well as other relations in the the space of form factors of scaling fields, have to be studied deeper, due to a possible appearance of the fields, satisfying the conditions (3)-(4). The following example illustrates, that we have to be very careful with an analysis of the off critical fields structure. We consider the multi particle form factors of the perturbed fields

$$\psi_k\bar{\psi}^{\dagger}_k = \Phi^{(N)}_{2k-N,N-2k}\,.$$

The expressions for these form factors, computed by a direct application of eq. (28), up to a normalization, coincide, as functions of rapidities, with the corresponding results for the field $\Phi^{(2)}_{2k,-2k} = \Phi^{(N-2)}_{2k-N,N-2k}$. The equality follows from the identities for trigonometric functions. In that way, after the perturbation, we get, formally, that two fields have the same form factors, while they are different at the criticality. This disagrees with the statement, that conformal and massive scaling fields should be in one-to-one correspondence [13]. The point is, that, this formal coincideness of expressions for different fields happens in a very specific case, in which the form factor prescription has to be worked out more carefully. Using the explicit values for the conformal dimensions (21), we see, that the following relations between the scaling dimensions of these fields takes place

$$2d^{(0)}_{2k} - 2d^{(2)}_{2k} = (2 - 2D_1). \tag{41}$$

Since $D_1 = \Delta_{\varepsilon_1}$ is the conformal dimension of the thermal operator ε_1, this equation is exactly the condition (3), that the fields $\psi_k \bar{\psi}^\dagger_k$ are in the first order resonance[4] with fields $\Phi^{(2)}_{2k,-2k}$. I.e. there is an ambiguity (4) in its definition

$$\psi_k \bar{\psi}^\dagger_k \sim \psi_k \bar{\psi}^\dagger_k + \mathrm{const}\, \lambda\, \Phi^{(2)}_{2k,-2k}\,. \tag{42}$$

Moreover, the spinless fields $\psi_k \bar{\psi}^\dagger_k$ formally have divergent expectation values, therefore, we need to work out a regularized prescription for its normalized multi particle matrix elements.

To introduce a well-defined form factor of the field $\psi_k \bar{\psi}^\dagger_k$, we propose to use a freedom in the choosing the index N. Namely, let us provide an analytic continuation of the expression for the multi-point form factor of this field, by changing $N \to N + \epsilon$, where the regularization parameter ϵ is a small number, which will be set to zero in the end. After the change, the vacuum expectation value becomes finite and proportional to $1/\epsilon$. We can divide the ϵ-deformed expressions, obtained from eq. (28), to the correspondent finite value of VEV and normalize the null point form factors to 1. The same analytic continuation and normalization are provided for the field $\Phi^{(2)}_{2k,-2k}$. Then, we propose to define the regularized form factors of $\psi_k \bar{\psi}^\dagger_k$ as a result of the formal calculation

$$\lim_{\epsilon\to 0} \frac{1}{\epsilon} \left(\frac{\langle \psi_k \bar{\psi}^\dagger_k | \{\beta\}, \{\beta'\} \rangle^{(\epsilon)}_{(n,n)}}{\langle \psi_k \bar{\psi}^\dagger_k \rangle^{(\epsilon)}} - \frac{\langle \Phi^{(2)}_{2k,-2k} | \{\beta\}, \{\beta'\} \rangle^{(\epsilon)}_{(n,n)}}{\langle \Phi^{(2)}_{2k,-2k} \rangle^{(\epsilon)}} \right). \tag{43}$$

This is one of the possible prescriptions in fixing the ambiguity (4). In the ultra violet regime, we should provide an analogous regularization for the field $\psi_k \bar{\psi}^\dagger_k$, which leads to vanishing expectation values of regularized fields.

To support this construction, we applied the regularization procedure for finding the form factors of the descendant field $E_1 = W^{(3)}_{-1} \bar{W}^{(3)}_{-1} \varepsilon_1$ in the $N = 4$ model.[5] The field E_1 in this theory satisfies the first order resonance condition with the identity operator. For form factors of this field, the regularization prescription provides a correct result, which can be calculated from another consideration. Namely, for general N, the form factors of the field E_1 have a very simple property. It can be expressed in terms of form factors of the first energy field in a following way

---

[4] The same is true for the Z_N charged fields $\psi_k \bar{\psi}_k$ and $\Phi^{(2)}_{2k,2k}$.

[5] This theory can also be considered as a well-known sine-Gordon model in the reflectionless point, with the parameter $\beta^2 = \frac{1}{3}$. The form factors of the exponential fields in this case can be extracted from the results of Refs. [22, 33]

$$\begin{aligned}\langle E_1|\{\beta\},\{\beta'\}\rangle_{(n,n)} = \frac{\sum\left(e^{2\beta_j}-e^{2\beta'_j}\right)}{\sum\left(e^{\beta_j}+e^{\beta'_j}\right)}\frac{\sum\left(e^{-2\beta_j}-e^{-2\beta'_j}\right)}{\sum\left(e^{-\beta_j}+e^{-\beta'_j}\right)}\times \\ \times\langle\varepsilon_1|\{\beta\},\{\beta'\}\rangle_{(n,n)}\,. \qquad (44)\end{aligned}$$

This equation comes from the fact, that in CFT [27] the energy field ε_1 has a null vector at the level 2. This means, that there is a linear relation between the descendants [19] $W^{(3)}_{-2}\varepsilon_1$ and $L_{-1}W^{(3)}_{-1}\varepsilon_1$. Now we have to take into account, that the modes $W^{(3)}_{-2}$ of the $W_3(z)$ current acts on the energy field ε_1 as the spin 2 integral of motion. Due to its even spin, it is odd with respect to the charge conjugation transformation, and have the form

$$\langle W^{(3)}_{-2}\varepsilon_1|\{\beta\},\{\beta'\}\rangle_{(n,n)} = \sum^{n}\left(e^{2\beta_j}-e^{2\beta'_j}\right)\langle\varepsilon_1|\{\beta\},\{\beta'\}\rangle_{(n,n)}\,. \qquad (45)$$

This completes the explanation of the eq. (44).

We suppose to study another relations, following from the null vector conditions, in a separate publication.

4.3 Explicit expressions for form factors

Finally, let us derive explicit expressions for the form factors, which will be used in the correlation functions studies. With the definitions (27)-(31), the equation (28) for the disorder fields has a conventional form

$$\begin{aligned}\langle\Phi^{(k)}_{m,-m}|\{\beta\},\{\beta'\}\rangle_{(n,n)} = \frac{(-1)^n}{(2\,\sin\frac{\pi}{N})^n}e^{\frac{m}{N}\sum(\beta_j-\beta'_j)}\times \\ \sum_{\{a_j,b_j\}}\prod_j a_j b_j e^{\frac{i\pi}{2N}((k+1)(a_j-b_j)-m(a_j+b_j))}\times \\ \langle\!\langle\mathcal{Z}_{a_1}(\beta_1)\cdots\mathcal{Z}^\dagger_{b_n}(\beta'_n)\rangle\!\rangle\langle\Phi^{(k)}_{m,-m}\rangle\,. \qquad (46)\end{aligned}$$

As an example, we write down the explicit expressions for the first form factors of the disorder operator

$$\begin{aligned}&\langle\mu_k|0\rangle = \langle\mu_k\rangle\,,\\ &\langle\mu_k|\beta_1,\beta'_1\rangle_{1^\dagger 1} = -\zeta^\dagger(\beta_1-\beta'_2)\langle\mu_k\rangle\{k\}e^{\frac{N-2k}{2N}(\beta_1-\beta'_1)}\,,\\ &\langle\mu_k|\beta_1,\beta_2,\beta'_1,\beta'_2\rangle_{1^\dagger 1^\dagger 11} = \zeta(\beta_{12})\zeta(\beta'_{12})\prod_{ij}^{2}\zeta^\dagger(\beta_i-\beta'_j)\\ &\quad\times\frac{1}{4}\langle\mu_k\rangle\{k\}^2 e^{\frac{k-N}{N}(\beta_1+\beta_2)-\frac{k}{N}(\beta'_1+\beta'_2)}\\ &\qquad\times\Big(\nu_1\tau_1+\frac{\{2\}\{k+1\}}{\{1\}\{k\}}\nu_2+\frac{\{2\}\{k-1\}}{\{1\}\{k\}}\tau_2\Big)\,, \qquad (47)\end{aligned}$$

where ν_j and τ_j are j-th symmetric polynomials of the variables e^{β_i} and $e^{\beta'_i}$, respectively, and we introduced the notation $\{a\} = \sin(\frac{\pi a}{N})$. In eqs. (47), we have written explicitly the normalization of scaling fields to its VEV [33]. The exact values of VEVs for the relevant for us fields [7] will be written below. It is rather direct task to find out expressions for the higher particle form factors. However, in our numerical computations, we will not need it.

Another simplest example is the case of parafermionic currents. Our candidate for the parafermionic current form factors can be easily found by using the equation (28)

$$
\begin{aligned}
&\langle\psi|\beta'_1\rangle_1 = C_\psi \; e^{\frac{N-1}{N}\beta'_1} \,, \\
&\langle\psi|\beta_1,\beta'_1,\beta'_2\rangle_{1^\dagger 11} = -C_\psi\{2\}\zeta(\beta'_{12})\prod_j^2 \zeta^\dagger(\beta_1-\beta'_j)\times \\
&\qquad\qquad\qquad\qquad \times e^{\frac{1}{N}\beta_1+\frac{N-2}{N}(\beta'_1+\beta'_2)} \,, \\
&\langle\psi|\beta_1,\beta_2,\beta'_1,\beta'_2,\beta'_3\rangle_{1^\dagger 1^\dagger 111} = \frac{1}{4}C_\psi\{2\}^2\zeta(\beta_{12})\prod_{i<j}^3\zeta(\beta'_{ij}) \\
&\qquad \times\prod_i^2\prod_j^3\zeta^\dagger(\beta_i-\beta'_j)e^{(\frac{1}{N}-\frac{1}{2})(\beta_1+\beta_2)-\frac{1}{N}(\beta'_1+\beta'_2+\beta'_3)} \\
&\qquad \times\Big(\tau_2+\nu_1\tau_1+\nu_2(1+2\cos\frac{2\pi}{N})\Big) \,, \qquad (48)
\end{aligned}
$$

etc. By C_ψ we denoted the normalization factor, determining the one particle form factor. We will fix it explicitly in eq. (71).

5 Correlation functions

In this section, we develop the conformal perturbation theory and compare long and short distance asymptotics of correlation functions of scaling fields. Our aim is to demonstrate, that the asymptotics are in agreement with each others at the intermediate distances, and, therefore, give an effective description of correlation functions at all scales. See Refs. [1, 40, 7, 31, 41] for other results in this direction.

Before proceed further, let us make an important comment. The long distance behaviors of correlation functions are described in terms of the mass of the lightest particle parameter. The exact relation between the mass and the coupling constant can be derived via TBA technique [2]. For the sine-Gordon model, this was done in the Al. Zamolodchikov's paper [3]. In our case the mass-coupling constant relation can be found from the results of Ref. [35]. To simplify expressions, here and below we use the notations:

$$
\gamma(a) = \frac{\Gamma(a)}{\Gamma(1-a)} \,, \quad u = \frac{1}{N+2} \,, \quad \kappa = M\frac{\Gamma(2/N)\Gamma(1-1/N)}{\Gamma(1/N)} \,. \quad (49)
$$

Applying these definitions, the explicit relation between the parameters M and

λ is given as following

$$(2\pi\lambda)^2 = \kappa^{4(1-2u)}\gamma(u)\gamma(3u)\,. \tag{50}$$

5.1 Conformal Perturbation Theory and exact VEVs

In this subsection, we consider the basic notions of the conformal perturbation theory [1] and give explicit values of the VEVs [7, 8] of the fields, which we will need in the computations.

In what follows, we develop the conformal perturbation theory [1] for the two point functions of scaling fields Φ_a and Φ_b

$$\langle\Phi_a(z,\bar{z})\Phi_b(0)\rangle = \sum_l C_{a,b}^{O_l}(z,\bar{z})\ \langle\, O_l(0)\,\rangle\,, \tag{51}$$

where $C_{a,b}^{O_l}(z,\bar{z})$ are the structure functions, and operators O_l form the basis in the space of scaling fields. Functions $C_{a,b}^{O_l}(z,\bar{z})$ can be expanded in the perturbation series

$$\begin{aligned} C_{a,b}^{O_l}(z,\bar{z}) = |z|^{2\Delta_{O_l}} z^{-\Delta_{\Phi_a}-\Delta_{\Phi_b}}\ \bar{z}^{-\bar{\Delta}_{\Phi_a}-\bar{\Delta}_{\Phi_b}} \times& \\ \times\left(C_{a,b}^{(0)O_l} + \lambda|z|^{2(1-D_1)}C_{a,b}^{(1)O_l} + \cdots\right)&, \end{aligned}$$

where the coefficients $C_{a,b}^{(0)O_l}$ are the structure constants from the conformal fields theory, while the first order corrections $C_{a,b}^{(1)O_l}$ can be expressed through the integrals of correlation functions in CFT

$$C_{a,b}^{(1)O_l} = \int d^2y\ \langle\Phi_a(0)\Phi_b(1)\varepsilon_1(y,\bar{y})O_l(\infty)\rangle\,. \tag{52}$$

The vacuum expectation values $\langle O_l\rangle$ of the scaling fields O_l, appearing in the equation (51), have a non-perturbative nature [1]. These fundamental quantities depend on the normalization prescription for the fields. We find VEVs, assuming that the scaling fields are satisfy the standard conformal normalization prescription

$$\langle O(z,\bar{z})O^{\dagger}(0)\rangle = |z|^{-4\Delta_O} + \cdots\,, \quad |z|\to 0\,. \tag{53}$$

The exact VEVs for physically important operators $\Phi_{m,\bar{m}}^{(k)}$ and $E_k = W_{-1}^{(3)}\overline{W}_{-1}^{(3)}\varepsilon_k$ in scaling Z_N Ising model were found in Refs. [7, 8], following the approach of Refs. [4, 35, 26]. The vacuum expectation values for the disorder fields are

$$\begin{aligned} \langle\mu_k\rangle = \kappa^{2d_k}\ \times& \\ \exp\int_0^{\infty}\frac{dt}{t}\left(\frac{\sinh kut\,\sinh(N-k)ut}{\sinh t\tanh Nut} - 2d_k e^{-2t}\right)&\,. \end{aligned} \tag{54}$$

It is interesting, that for even values of k, the integral above can be calculated, giving

$$\langle\mu_{2k}\rangle = \kappa^{2d_{2k}}\left(\frac{\gamma(\frac{1}{N+2})}{\gamma(\frac{2k+1}{N+2})}\right)^{1/2}\left(\frac{N+2}{N}\right)^{\frac{k(N-2k)}{N}}\prod_{i=1}^{k}\frac{\gamma(\frac{2i-1}{N})}{\gamma(\frac{2i-1}{N+2})}\,. \tag{55}$$

For a general case, $\Phi^{(k)}_{m,-m} = (\psi^\dagger)^l(\overline{\psi})^l\mu_k$, where $m = k-2l$, the vacuum expectation values of fields are elegantly expressed in terms of $\langle\mu_k\rangle$ as

$$\frac{\langle\Phi^{(k)}_{m,-m}\rangle}{\langle\mu_k\rangle} = \left(\frac{\kappa(N+2)}{N}\right)^{\frac{k^2-m^2}{2N}}\prod_{i=0}^{l-1}\frac{(i+1)\gamma(\frac{i+1}{N})}{(k-i)\gamma(\frac{k-i}{N})}\,. \tag{56}$$

In particular, VEVs of thermal fields $\varepsilon_k = (\psi)^k(\overline{\psi})^k\mu_{2k}$ correspond to the $m=0$ case in the equation above:

$$\begin{aligned}\langle\varepsilon_k\rangle = \quad & \kappa^{2D_k}\left(\frac{N+2}{N}\right)^k\frac{(k!)^2}{(2k)!}\left(\frac{\gamma((2k+1)u)}{\gamma(u)}\right)^{\frac{1}{2}}\times \\ & \prod_{i=1}^{k}\frac{\gamma^2(\frac{i}{N})}{\gamma(\frac{2i}{N})\gamma((2i+1)u)}\,.\end{aligned} \tag{57}$$

We also obtained the exact result for the expectation values of the normalized descendent fields $E_k = W^{(3)}_{-1}\overline{W}^{(3)}_{-1}\varepsilon_k$

$$\begin{aligned}\frac{\langle E_k\rangle}{\langle\varepsilon_k\rangle} = \quad & \kappa^2\frac{(N+2)^2}{2N}\frac{\Gamma^2(1+\frac{k+1}{N})\Gamma^2(1-\frac{k}{N})}{\Gamma^2(1-\frac{k+1}{N})\Gamma^2(1+\frac{k}{N})}\times \\ & \frac{\Gamma(1+\frac{2k}{N})\Gamma(1-\frac{2k+2}{N})}{\Gamma(2-\frac{2k}{N})\Gamma(2+\frac{2k+2}{N})}\,.\end{aligned} \tag{58}$$

5.2 Correlation functions $\langle\mu_1(x)\mu_1(0)\rangle$

In our previous paper [7], we studied long and short distance asymptotics of the correlation functions $\langle\sigma_1(x)\sigma_1^\dagger(0)\rangle$ and $\langle\mu_1(x)\mu_1^\dagger(0)\rangle$ of order and disorder fields. For completeness, we collect the correspondent ultra violet expansion data in the appendix A. We found, that IR and UV asymptotics of the correlators match at the intermediate distances. This confirm our expressions for VEVs, as well as form factor expressions.

In this subsection, we want to discuss a more complicate case of the correlation function, including two disorder fields $\langle\mu_1(x)\mu_1(0)\rangle$. We found, that studying this case is rather instructive, since it gives an example, where the resonance fields (3)-(4) appear at the short distance expansion for all integer $N = 2, 3, \ldots$

The long distance expansion of this correlation function is provided in a standard way. In a two particle form factor approximation, we have the spectral decomposition

$$\begin{aligned}&\langle\mu_1(z,\bar z)\mu_1(0)\rangle = \langle\mu_1\rangle^2 + \\ &+\int d\theta_1\ d\theta_2\Big(\langle\mu(z,\bar z)|\theta_1,\theta_2\rangle_{1,1^\dagger}\ \langle\mu^\dagger(0)|\theta_2+i\pi,\theta_1+i\pi\rangle_{1,1^\dagger}+ \\ &\qquad +\langle\mu(z,\bar z)|\theta_1,\theta_2\rangle_{1^\dagger,1}\ \langle\mu^\dagger(0)|\theta_2+i\pi,\theta_1+i\pi\rangle_{1^\dagger,1}\Big)+\cdots\,. \end{aligned}\tag{59}$$

Studying the short distance behaviors is more involved. Indeed, we have the following leading terms in the conformal perturbation theory expansion (51):

$$\begin{aligned}\langle\mu_1(z,\bar z)\mu_1(0)\rangle &= C^{\mu_2}_{\mu_1\mu_1}(r)\langle\mu_2\rangle + C^{\eta}_{\mu_1\mu_1}(r)\langle\eta\rangle + \\ &+C^{\psi\bar\psi}_{\mu_1\mu_1}(r)\langle\psi\bar\psi^\dagger\rangle+\cdots\,,\qquad |z|=r\,.\end{aligned}\tag{60}$$

Where we introduced the notation η for the field $\eta=\Phi^{(2)}_{4,2}$. The leading contribution to the correlator (60) at the short distances comes from the zero order term $C^{(0)\mu_2}_{\mu_1\mu_1}(r)\langle\mu_2\rangle$ of the perturbation theory. It can be computed using CFT structure constants, found in Ref. [9], and the exact vacuum expectation value for the second disorder field μ_2 (54).

$$C^{(0)\mu_2}_{\mu_1\mu_1}(r)\ \langle\mu_2\rangle = r^{-2d_1}\frac{\gamma\left(\frac{1}{N}\right)}{\gamma(2u)}(r\kappa)^{2\frac{N-2}{N(N+2)}}\left(\frac{N+2}{N}\right)^{\frac{N-2}{N}}\,,\qquad u=\frac{1}{N+2}\,.\tag{61}$$

At the first order perturbation theory we have the term, including the field η. The contribution from this field to the perturbative expansion can be expressed in terms of the product of the CFT structure constants $C^{(0)\mu_2}_{\mu_1\mu_1}\ C^{(0)\eta}_{\mu_2\varepsilon_1}$, multiplied by the simple two dimensional integral over y from the correlation function

$$\langle\mu_1(0)\mu_1(1)\varepsilon_1(y,\bar y)\eta(\infty)\rangle_{CFT} = |y|^{2u}|1-y|^{2u}\,.$$

Using the exact result (56) for VEV of the field $\eta=\Phi^{(4)}_{2,-2}$, we obtain the analytic result for this term in the first order approximation

$$C^{\eta}_{\mu_1\mu_1}(r)\langle\eta\rangle = \frac{r^{-2d_1}}{16}(r\kappa)^{2\frac{(N+2)^2-9}{N(N+2)}}\frac{\gamma\left(\frac{1}{N}\right)\gamma\left(\frac{3}{N}\right)\gamma^3(u)\,u^2}{(1+2u)^2\gamma(3u)\gamma^2(2u)}\left(\frac{1}{Nu}\right)^{2\frac{N-1}{N}}\,.\tag{62}$$

A more difficult task is to compute the first order correction to the structure function $C^{\mu_2}_{\mu_1\mu_1}(r)$ and the contribution, coming from the term $\psi\bar\psi^\dagger$. We note that the conformal dimensions of the fields $\psi\bar\psi^\dagger$ and μ_2 satisfy to the relation

$$2\Delta_\psi - 2\Delta_{\mu_2} = 2\frac{N}{N+2} = 2\left(1-\Delta_{\varepsilon_1}\right)\,.\tag{63}$$

The first order resonance condition (3) appears now at the short distance expansion. The equation (63) basically means, that two terms in the perturbation theory will have the same powers of r. Whenever this happens, we expect, that divergencies coming from the contributions from the term $C^{\psi\bar\psi^\dagger}_{\mu_1\mu_1}(r)\langle\psi\bar\psi^\dagger\rangle$

should be canceled by divergencies, appearing from the first order correction to the structure function $C^{(1)\mu_2}_{\mu_1\mu_1}(r)\langle\mu_2\rangle$. This phenomena, in general, should lead to logarithmic terms in a perturbative expansion.

The appropriate contributions to the resonance terms can be computed within the Z_N Ising model settings by providing the analytic continuation in the parameter N, as it was discussed before. I.e., we can compute the contributions after the change $N \to N + \epsilon$, and then consider the limit $\epsilon \to 0$. To check the validity of this approach, we also perform the computations in a different method. We use the fact, that the parafermionic CFT with the central charge (1) belongs to the series of W algebra symmetric unitary minimal CFT $\mathcal{W}M^{(p)}_N$, where the parameter p is chosen to be $p = N + 1$ [27]. The field theory (2), from the view point of this model, is the simplest representative of the series of integrable perturbations of $\mathcal{W}M^{(p)}_N$ minimal models by the primary field with the conformal dimension $D^{(p)}_1 = 1 - \frac{N}{p+1}$, which, in our case, coincides with the first energy operator. To study contributions of the resonance fields we can do the deformation of the $\mathcal{W}M^{(p)}_N$ by changing the parameter p in such a way that the variable $u = 1/(p + 1)$ would have the form

$$u_\epsilon = \frac{1}{N+2} - \epsilon\,.$$

For small non-zero ϵ the expressions for the resonance fields contributions are finite and well-defined. The result of computations can be written in the following way

$$\begin{aligned}
&C^{(1)\mu_2}_{\mu_1\mu_1}(r)\langle\mu_2\rangle = r^{-2d_1}(r\kappa)^{\frac{2}{N}(N-2+(N+2)u_\epsilon)}\frac{2}{N^2}\frac{(1-(N+1)u_\epsilon)^2}{(1-Nu_\epsilon))^2}\times\\
&\times\frac{\gamma\left(\frac{1}{N}\right)\gamma\left(Nu_\epsilon\right)}{\gamma\left(2u_\epsilon\right)}\frac{\gamma\left(u_\epsilon\right)\gamma^2\left(\frac{1+Nu_\epsilon}{2}\right)\gamma^2\left(\frac{1-(N-2)u_\epsilon}{2}\right)\gamma\left(\frac{1-(N+2)u_\epsilon}{2}\right)}{\gamma((N+1)u_\epsilon)(Nu_\epsilon)^{\frac{N-2}{N}}}\times\\
&\times\left(1+\epsilon(N+2)\chi_0+O(\epsilon^2)\right)\,,
\end{aligned}\tag{64}$$

To simplify the resulting expression, we introduced a shorthand notation for the term in the last line

$$\begin{aligned}
\chi_0 = -\frac{1}{2}\Bigg(&\psi\left(\frac{1}{2}+\frac{(N-2)u}{2}\right)+\psi\left(\frac{1}{2}-\frac{(N-2)u}{2}\right)-\frac{2}{N}\gamma_E+\\
&+(N+2)\log\left(\frac{N+2}{N}\right)\Bigg)\,.
\end{aligned}$$

The symbol γ_E is reserved for the Euler's constant and ψ stands for the logarithmic derivative of the gamma function $\psi(x) = \Gamma'(x)/\Gamma(x)$. (Unfortunately, this standard notation conflicts with the usual symbol, which we choose for the parafermionic current $\psi(z)$).

The contribution, which comes from the term proportional to $\langle\psi\bar{\psi}^\dagger\rangle$, is found to be of a similar form

$$C^{\psi\bar{\psi}^\dagger}_{\mu_1\mu_1}(r)\langle\psi\bar{\psi}^\dagger\rangle = r^{-2d_1}(r\kappa)^{2\frac{N-2}{N}(2-(N+2)u_\epsilon)}\frac{(1-(N+1)u_\epsilon)^2}{(1-2u_\epsilon)^2}\times$$

$$\begin{aligned}&\times\left(\frac{1}{Nu_\epsilon}\right)^{2\frac{N-1}{N}}\frac{\gamma\left(Nu_\epsilon\right)}{\gamma\left(u_\epsilon\right)}\gamma\left(\frac{1-Nu_\epsilon}{2}\right)\gamma\left(\frac{1-(N-2)u_\epsilon}{2}\right)\times\\&\times\gamma\left(\frac{1}{Nu_\epsilon}-1-\frac{1}{N}\right)\gamma\left(\frac{N+2}{N}-\frac{1}{Nu_\epsilon}\right)\times\\&\times(1+\epsilon(N+2)\chi_1+O(\epsilon^2))\,,\end{aligned}\tag{65}$$

where

$$\chi_1=-\frac{1}{2}\left(\psi\left(\frac{1}{2}+\frac{Nu}{2}\right)-\psi\left(\frac{1}{2}-\frac{Nu}{2}\right)\right)+\chi_0\,.$$

Now we look for the regularized expressions in the limit $\epsilon\to 0$.

It can be derived from the equations (64)-(65), that both terms have a single pole at $\epsilon=0$. However, as we expected from a general consideration, the sum of residues at this pole is zero, and the total contribution of these resonance terms to the correlation functions is well defined in the $\epsilon\to 0$ limit. Computing the limit, we find, that the correctly defined sum of two resonance terms, is given as

$$\begin{aligned}&\left[C^{(1)\mu_2}_{\mu_1\mu_1}(r)\langle\mu_2\rangle+C^{\psi\bar{\psi}^\dagger}_{\mu_1\mu_1}(r)\langle\psi\bar{\psi}^\dagger\rangle\right]^{\mathrm{reg}}=\frac{\gamma(\frac{1}{N})}{2N^2}r^{-2d_1}(r\kappa)^{2\frac{N-1}{N}}\left(\frac{N+2}{N}\right)^{\frac{N-2}{N}}\\&\times\Bigg(8(\gamma_E-1)+4N\log r\kappa e^{\frac{3}{2}\gamma_E-1}+2(N+2)\left(\psi(\frac{1}{N})+\psi(-\frac{1}{N})\right)-\\&\qquad-N\left(\psi(\frac{2}{N+2})+\psi(\frac{N}{N+2})\right)\Bigg)\,.\end{aligned}\tag{66}$$

The numerical data for long and short distance asymptotic expansions are shown in the Fig. 5 for the case N=7.



Fig. 5. Correlation function of disorder fields at N=7.

The dashed curve is for UV asymptotics, while the full line denotes the form factor decomposition, up to two particles. We observe, that there is a matching of the asymptotics at the intermediate distances. We found, that, in the region $0.01\leq Mr\leq 1$, the long and short distance asymptotics agree with the relative

error around 1 percent. This, rather good, numerical preciseness confirms our hypothesis on the identification of form factors, as well as the short distance regularization prescription.
For other values of the parameter N, we provided similar numerical computations. We found, that the relative error decreases with increasing of the number N. For large N the error becomes smaller. For example, for N=11 case, it is already less than 0.1 percent. The agreement between data can be further improved by taking into account higher particles form factors, however this is beyond the scope of the present paper.

In the limit of large parameters N we find, that our short distance expansion function behaves as

$$\frac{\langle\mu_1(z,\bar{z})\mu_1(0)\rangle}{\langle\mu_1\rangle^2} = 1 + \frac{1}{N^2}\left(-2\Omega + (\Omega^2 - 4\Omega + \frac{9}{2})M^2r^2\right) + \cdots\,, \quad (67)$$

where $\Omega = \log\left(\frac{Mre^{\gamma_E+1}}{2}\right)$. We checked, using the Mellin transform, that our form factor expression leads to the same expansion at the small distances up to the order N^{-3} . In the next orders new corrections can appear from the higher particle form factors.

Another non-trivial test for the correctness of our expressions is the limit to the Ising model point $N = 2$. In the case, when $N \to 2$, our ultra violet asymptotics leads to the following answer

$$\langle\mu_1(z,\bar{z})\mu_1(0)\rangle = \frac{1}{r^{\frac{1}{4}}}\left(1 - \frac{Mr}{2}\log\frac{Mre^{\gamma_E}}{8} + \cdots\right)\,, \quad (68)$$

which agree with the known short distance expansion for the Ising model disorder parameter correlation function [15, 36].

5.3 Correlation functions of parafermionic currents

In this subsection we would like to compare short and long distance asymptotics for the two point correlation functions of parafermionic currents ψ and $\psi^\dagger$ at the region $Mr \sim 1$. We think, that a consideration of this case might be interesting. First of all, because the correlator of free fermions is one of the most simplest in the Ising model. Unlike the case of order and disorder fields, it is given exactly by the Bessel function, i.e. by the solution of a linear equation, which is simplier, than the Painleve equation [15, 36]. From the other side, this is one of few examples of correlation functions of operators with fractional spins.

We recall, that in the unperturbed CFT, the currents $\psi(z)$ and $(\psi^\dagger)$ have conformal dimensions $(\Delta_1, 0)$ and $(\Delta_{N-1}, 0)$, defined by eq. (12). In the ultra violet region the correlation function of these fields is expected to have a form (see eq. (18))

$$\langle\psi(z,\bar{z})\psi^\dagger(0)\rangle = z^{-2\Delta_1} + \cdots\,, \quad |z| \to 0\,, \quad (69)$$

which is natural definition of the conformal normalization in the given case of operators with non-trivial spins.

5.3.1 General N case

The leading contribution to the infra red asymptotics of the correlation function of parafermionic currents comes from the one particle form factor approximation. We will see, that it already leads to the nice agreement between the asymptotics at the intermediate distances. Using the integral representation for the modified Bessel function, we can express the one particle contribution in the following analytic form

$$\langle\psi(z,\bar{z})\psi^{\dagger}(0)\rangle = 2\; C_{\psi}^{2} K_{\frac{N-1}{N}}(r) + \cdots\,, \qquad z = r\,. \tag{70}$$

For simplicity, we choose here and below the space coordinate to be zero, which corresponds to the real $z = r$.

In the equation (70) we took into account, that for the fields with spin the vacuum expectation values vanishes. Our proposal for the exact value of the multiple C_{ψ}^{2}, leading to the conformal normalization (69) of parafermionic currents, is

$$C_{\psi}^{2} = \frac{\Gamma(1+\frac{1}{N})}{\Gamma(1-\frac{1}{N})}\left(\frac{N+2}{N}\right)^{2\frac{N-1}{N}} \kappa^{2\frac{N-1}{N}}\; S_2^2(2\pi + \frac{2\pi}{N})\,. \tag{71}$$

The constant $S_2(2\pi + \frac{2\pi}{N})$ is defined by eq. (90) in the appendix B. This expression can be obtained, following the ideas of Ref. [34]. It comes naturally from analysis of the divergency in the VEV of the operator $\psi\bar{\psi}^{\dagger}$, which can be effectively provided in the framework of the W-symmetric CFT, perturbed by adjoint field, by use of the results of the paper [35]. In another way, it can be obtained by analyzing the deformed parafermionic currents normalization, see Ref. [8]. Further, we will see, that this normalization coincides with the known exact results for $N = 2$ and $N \to \infty$ cases. We also establish numerical checks, by matching the long and short distance asymptotics for correlators for arbitrary N.

From the general arguments of the conformal perturbation theory [1], we found, that the short distance expansion of the correlator of parafermionic currents has the form

$$\begin{aligned}\langle\psi(z,\bar{z})\psi^{\dagger}(0)\rangle = \frac{1}{z^{2\Delta_1}}\Big(&1 + (\kappa r)^{2}A_1 + (\kappa r)^{\frac{4N}{N+2}}A_2 + \\ &+(\kappa r)^{\frac{4(N+3)}{N+2}}A_3 + (\kappa r)^{\frac{4N}{N+2}+2}A_4 + \\ &+(\kappa r)^{\frac{4(N+1)}{N+2}+2}A_5 + (\kappa r)^{\frac{6(N+4)}{N+2}}A_6 + \cdots\Big)\,.\end{aligned} \tag{72}$$

Here and below in this subsection, we use in the short distance expansions the notation $r = |z|$. For the computation of the coefficients $A_j,\ (j = 1,\ldots,6)$ in this equation, it is convenient to use the quantum equations of motion (37)

$$\frac{\partial}{\partial\bar{z}}\psi(z,\bar{z}) = \lambda\sqrt{\frac{2}{N}}\,\Phi^{(2)}_{2,0}(z,\bar{z})\,. \tag{73}$$

We recall, that the fields $\Phi^{(2)}_{2,0} = (\psi)\epsilon_1$ and $\Phi^{(2)}_{-2,0} = (\psi^\dagger)\epsilon_1$, have different left and right conformal dimensions $(\Delta_{\mu_2}, \Delta_{\epsilon_1})$ defined by eqs. (7), (23).

To define the coefficient A_1, we integrate over $\bar{z}$ the first term in the decomposition of the following two point correlation function

$$\langle \bar{\partial}\psi(z,\bar{z})\psi^\dagger(\zeta,\bar{\zeta})\rangle = \lambda\sqrt{\frac{2}{N}}\langle \Phi^{(2)}_{2,0}(z,\bar{z})\psi^\dagger(\zeta,\bar{\zeta})\rangle\,. \tag{74}$$

In that prescription the coefficient A_1 is computed in a simple way from the CFT three point correlation function

$$\langle \Phi^{(2)}_{2,0}(0)\psi^\dagger(1)\varepsilon_1(\infty)\rangle_{CFT}\,,$$

multiplied to the expectation value of the field ε_1. An effective method of computation of other coefficients in the equation (72) is to integrate twice the series expansion for the following correlation function

$$\langle \bar{\partial}\psi(z,\bar{z})\bar{\partial}\psi^\dagger(\zeta,\bar{\zeta})\rangle = \frac{2\lambda^2}{N}\langle \Phi^{(2)}_{2,0}(z,\bar{z})\Phi^{(2)}_{-2,0}(\zeta,\bar{\zeta})\rangle\,. \tag{75}$$

More explicitly, starting from the conformal perturbation theory expansion for the two point correlator

$$\begin{aligned} &\langle \Phi^{(2)}_{2,0}(z,\bar{z})\Phi^{(2)}_{-2,0}(\zeta,\bar{\zeta})\rangle = \frac{|z-\zeta|^{\frac{4N}{N+2}}}{(z-\zeta)^{2\Delta_1}(\bar{z}-\bar{\zeta})^2}\times \\ &\Big(C^I(r) + C^{\epsilon_1}(r)\langle\epsilon_1\rangle + C^{\epsilon_2}(r)\langle\epsilon_2\rangle + C^{E_1}(r)\langle E_1\rangle + C^{\epsilon_3}(r)\langle\epsilon_3\rangle + \cdots\Big)\,, \end{aligned} \tag{76}$$

we obtain, by integration over $\bar{z}$ and $\bar{\zeta}$, the coefficient A_2 from the coefficient C^I. In a similar manner, the coefficients A_3, A_5 and A_6 are related, correspondingly, with the zeroth order structure functions $C^{\varepsilon_2}\langle\varepsilon_2\rangle$, $C^{E_1}\langle E_1\rangle$ and $C^{\varepsilon_3}\langle\varepsilon_3\rangle$. The coefficient A_4 is related with the first order correction term $C^{(1)\varepsilon_1}\langle\varepsilon_1\rangle$.

Using the free field realization for conformal primary fields [9, 11], one can find an integral representation for the leading corrections to the structure functions in the expansion (75). The integrals in the first order perturbation theory can be taken by applying the technique from Refs. [37]. The exact vacuum expectation values for the energy fields $\langle\epsilon_k\rangle$, $(k=1,2,3)$ and E_1 are given in eq. (57) and (58) respectively. The results for numerical coefficients A_k can be written as

$$\begin{aligned} A_1 &= -\frac{N+2}{2N^2}\frac{\gamma(\frac{1}{N})^2}{\gamma(\frac{2}{N})}\,, \\ A_2 &= \frac{(N+2)^2}{4N^2(N-2)}\gamma(u)\gamma(3u)\,, \\ A_3 &= -\frac{1}{72}\frac{(N+2)^4}{(N+3)(N+4)N^3}\frac{\gamma(u)}{\gamma(3u)}\frac{\gamma^2(\frac{1}{N})\gamma(\frac{2}{N})}{\gamma(\frac{4}{N})}\,, \\ A_4 &= -\frac{(N+2)^3}{16N^3(3N+2)}\frac{\gamma^2(\frac{1}{N})}{\gamma(\frac{2}{N})}\gamma(u)\gamma(3u)\;J_N\,, \end{aligned}$$

$$
\begin{aligned}
A_5 &= -\frac{(N+2)^5(N-4)}{8(N+1)(3N+4)}\,\frac{1}{(N+4)^2(N-2)^2N}\,\frac{\gamma^2(\frac{2}{N})}{\gamma(\frac{4}{N})}\,\frac{\gamma(4u)\gamma^2(u)}{\gamma^2(2u)}\,,\\
A_6 &= -\frac{1}{2400}\,\frac{(N+2)^5}{N^4(N+4)(N+5)(N+6)^2}\,\frac{\gamma^2(\frac{1}{N})\gamma^2(\frac{2}{N})\gamma^2(\frac{3}{N})}{\gamma(\frac{2}{N})\gamma(\frac{4}{N})\gamma(\frac{6}{N})}\times\\
&\qquad\qquad \frac{\gamma^2(2u)\gamma^2(u)}{\gamma(3u)\gamma(4u)\gamma(5u)}\,.
\end{aligned}
\tag{77}
$$

Note, that the coefficient A_4 was computed analytically only for particular values of N. This is because the multiple J_N in this coefficient represents the contribution, which is given by the rather complicate integral over the plane, from the CFT correlation function

$$
J_N = \frac{1}{\pi}\int d^2y\ \langle \Phi^{(2)}_{2,0}(1)\Phi^{(2)}_{-2,0}(0)\epsilon_1(y,\bar{y})\epsilon_1(\infty)\rangle_{CFT}\,. \tag{78}
$$

The asymptotics of the integral J_N, considered as a function of N, will be given for large values of N further, in eq. (84).

In the next subsection, we discuss the particular cases of N, for which the correlation function have specific features, including resonances.

5.3.2 Ising model N=2 case

We consider, first, the consistency of our formulae for N=2 with the known results from the Ising model. According to the fusion rules in Ising model case, there is no higher energy fields in the short distance expansion. So, we have to restrict ourself to the two first terms coming with the coefficients A_1 and A_2. A formal substitution $N=2$ leads to a divergency which is, of course, expected from our previous consideration of the resonance fields. When we consider the limit $N\to 2$, we find, that the residues at the pole $\frac{1}{N-2}$ are canceled, and the resulting expansion for correlator has a logarithmic scaling. Let us assume, that $z=r$ is chosen to be real. Then, the $N\to 2$ limiting value for correlator will have the form

$$
\begin{aligned}
\langle\psi(z,\bar{z})\psi(0)\rangle = \lim_{N\to 2} r^{-2\frac{N-1}{N}}\left(1+(\kappa r)^2A_1+(\kappa r)^{\frac{4N}{N+2}}A_2+\cdots\right) =\\
=\frac{1}{r}\left(1+\frac{(Mr)^2}{2}\log\left(\frac{Mr}{2}e^{\gamma_E-\frac{1}{2}}\right)+\cdots\right)\,.
\end{aligned}
\tag{79}
$$

This equation agrees with the form factor ultra violet expansion, where the exact correlator is given in terms of the modified Bessel function $K_1(r)$. In the numerical computations, we can formally put N in the general coefficients A_1 and A_2 to be close to 2. Then, the long and short distance expressions would match at the intermediate distances, as it is expected. Since the infra red expansion result is exact one, it is instructive to see for our short distance formulae and efficiency of the analytic continuation prescription. For example, the curves, depicted at Fig. 6. corresponds to N=2.00001.



Fig. 6. Parafermionic correlators at N=2.00001.

In the figure the dashed lines is given by the first three terms in the UV decomposition. The full line, here and below, represents the one particle form factor expansion. The dependence on N, in a vicinity of the Ising model case value of $N = 2$, is smooth. Matching between long and short distance asymptotics of the Ising model case serves as one of the confirmation, supporting the normalization constant C_ψ^2 (71).

5.3.3 Three states Potts model N=3 case

The model for $N = 3$ coincides with the scaling 3-state Potts model. Up to the normalization factors, the exact form factors for parafermionic currents (as well as for other primaries) were computed for this model in Ref. [29]. The correlation functions of order and disorder fields in this model were also studied in Refs. [7, 31].

According to the fusion rules for N=3 case, there is no energy fields ε_2 and ε_3 and we have to omit terms with the coefficients A_3, A_6. We show at Fig. 7, that even first three terms in the short distance expansion lead to a rather good agreement between correlation functions asymptotics.



Fig. 7. Parafermionic correlators at N=3.

In this case we can advance in the exact computations further. In particular, the coefficient A_4 can be calculated exactly, since the integral J_3 can be taken analytically as

$$J_3 = -\frac{9\sqrt{5}\Gamma^5(\frac{3}{5})\Gamma^4(\frac{1}{5})}{16\pi\Gamma^3(\frac{2}{5})\Gamma^2(-\frac{1}{5})} \,. \tag{80}$$

Then, the contribution from this first order perturbation theory term is given by the expression

$$(z\kappa)^{2+4\frac{N}{N+2}} A_4 \Big|_{N=3,z=r} = \frac{5}{3\cdot 11\cdot 2^6}\frac{\gamma^2(\frac{1}{3})\gamma^4(\frac{1}{5})}{\gamma(\frac{2}{3})\gamma^5(\frac{2}{5})}(r\kappa)^{\frac{22}{5}} \,. \tag{81}$$

It is possible also to compute the second order perturbation correction to the identity operator in the expansion (72), as well as the contribution of the descendant field $T\bar{T}$, those VEVs can be computed from the results of Refs. [38, 39]. With that corrections, and with the term with the coefficient A_5, taken into account, the long and short distance expressions agree up to the distances Mr=3.

5.3.4 N=4 model

The model for $N = 4$ describes a particular case of the Ahskin-Teller model. For the parafermionic fields, proceeding as before, we find, that the resonances appear at the short distance expansion, as well.

Namely, providing an analytic continuation of general expressions with parameter N, we can easily see, that the term with the coefficient A_3 diverges at $N \to 4$. However, the divergency, coming from this term, is canceled by the singular part of the term with the coefficient A_4, since the integral J_N has the following expansion, when N approaches 4

$$J_N = \frac{4}{3}\frac{1}{N-4} + \frac{1}{18} + O(N-4) \,. \tag{82}$$

In this case we meet the resonance condition (3) between the operators ϵ_2 and ϵ_1. As in the Ising model, let us look for an analytic continuation in N of the general expression. Finding the regular, in the parameter N, expansion, we obtain

$$\begin{aligned} &\lim_{N\to 4}\left((\kappa r)^{4\frac{N+3}{N+2}} A_3 + (\kappa r)^{2+4\frac{N}{N+2}} A_4\right) = \\ &\quad = \frac{3}{2^6\cdot 7}\frac{\gamma\left(\frac{1}{6}\right)}{\gamma^2\left(\frac{3}{4}\right)}(r\kappa)^{\frac{14}{3}}\log\left(\frac{r\kappa}{2}e^{\gamma^E - \frac{27}{28}}\right) \,. \end{aligned} \tag{83}$$

Again, the presence in the expansion fields, which are in the resonance, leads to logarithmic dependence of the short distance approximation. Let us note, that there is no divergency at the term A_5

$$\lim_{N\to 4}(\kappa r)^{2+4Nu+4u} A_5 = -\frac{3^{\frac{1}{2}}\cdot 3^5}{2^{\frac{1}{3}}\cdot 2^{10}\cdot 5\pi}\Gamma^2(\frac{1}{3})(\kappa r)^{\frac{16}{3}} \,.$$

Substituting that regularized expressions into the short distance expansion, we get a well defined expansion. The numerical data (where the contribution from the second order correction was taken into account), are shown at Fig. 8.



Fig. 8. Parafermionic correlators at N=4.

The relative error between asymptotic values in the region $0.0001 < Mr < 1$ in our numerical computations is less than 1 percent.

5.3.5 Large N case

The nice feature of the Z_N models is that the correlation functions in the ultra violet and infra red regions (72) are in agreement for an infinite set of models including those with an arbitrary integer $N \geq 2$. We propose in the large N limit the following approximation for the multiple J_N in eq. (77):

$$J_N = \frac{18}{5N}\left(1 + \frac{32}{3N^2} + O(\frac{1}{N^3})\right) . \tag{84}$$

Taking the A_4 term into account, we have, for the asymptotic expansion for correlation functions at $N \to \infty$ in the vicinity of $r = 0$, the following expansion

$$M^{-2\Delta_1}\langle\psi(z,\bar{z})\psi^\dagger(0)\rangle\Big|_{z=r} = \frac{1}{r^2}\left(T_0(Mr) + \frac{1}{N}\,T_1(Mr) + \cdots\right) . \tag{85}$$

Here, the functions $T_0(x)$ and $T_1(x)$ are given by the expressions, which agrees with the one particle form factors formulae for large N and small scales

$$\begin{aligned} T_0(x) &= 1 - \frac{x^2}{4} - \frac{x^4}{16}\left(\log x + \gamma_E - \frac{3}{4}\right) , \\ T_1(x) &= 2\log x - \frac{x^2}{2}(1 + \log x) + \\ &+ \frac{x^4}{16}\left(2\left(\log\frac{e^\gamma}{2} - \frac{7}{4}\right)\log x + 2\log^2\frac{e^\gamma}{2} - 5\log\frac{e^\gamma}{2} + \frac{13}{4}\right) . \end{aligned} \tag{86}$$

With that conventions, both, short and large, distance expansions produce similar looking curves for $N > 6$. The example of the correlation functions for the case $N = 11$ is given in the Fig. 9.



Fig. 9. Parafermionic correlators at N=11.

To show the difference between the asymptotics, we draw also the re-scaled correlators $r^{2\frac{N-1}{N}}\langle\psi\psi^\dagger\rangle$ in the Fig. 10.



Fig. 10. Re-scaled parafermionic correlators at N=11.

Let us note, that the relative error for ultra-violet and infra-red asymptotics at the region $0.000001 < Mr < 1$ become smaller with increasing of the number N. For example, for $N = 7$ model, the error in this region is 0.8 percent, while for $N = 20$ it is already 0.08 percent.

To deal with that small N cases we considered the A_4 term more accurately. We provided the Pade approximation for the integral J_N (78) between the known and fixed points $N = 4$ and $N = \infty$. The details of the computations are collected in the appendix C. Using this, we found, that the matching between the asymptotics of the correlation functions for small N is within 1 per cent at the distances $0.000001 < Mr < 1$.

6 Concluding remarks

In this paper we studied correlation functions of disorder fields and parafermionic currents for scaling Z_N Ising models and found that the long and short distance asymptotics approach each others at the intermediate distances. From one side, a matching between ultra violet and infra red asymptotics gives an effective way of studying basic behaviors of the correlation function of the theory at all distances. From the other side, it confirms our construction of the form factors of the scaling fields.

We discussed algebraic relations in the space of form factors and outline the set of problems related with the form factors of the descendant fields. [6] In particular, we stress the role of quantum equations of motion, by showing, that they appear naturally in the form factor prescription. From the other side, we demonstrated, that the equations of motion can be a very powerful tool in studying the ultra violet asymptotics, within the conformal perturbation theory. Their application allows to find coefficients in the short distance asymptotics, which are rather difficult to study by direct methods. Another useful method, which we applied in the analysis of form factors and conformal perturbation theory, is the W extended symmetry of the model. The W symmetric models of CFT and their integrable perturbations are, in general, rather complicate and we found that it is interesting, that correlation functions can be effectively studied for such models, at least, in the particular cases. We also stress the necessity of more deeper understanding the fields, which satisfy the resonance condition. We have shown, that these fields have unusual properties in both, form factor approach, as well as in the conformal perturbation theory, and have to be analyzed carefully.

In this paper, we did not consider ultra violet expansions for correlation functions of "heavy" fields O_a, which are in resonance with some field O_b (3)-(4). If we use the proposed form factor regularization prescription for these fields, then the calculation of the short distance behaviors for correlators become more subtle problem, and terms, including $\log^2 r$, will appear in the short distance expansion. The well-known example for this property is the correlation function $\langle\psi\bar{\psi}(x)\psi\bar{\psi}(0)\rangle$ in the Ising model.

Another interesting phenomena in studying of resonance fields is that, sometimes, the renormalization of primary fields is finite and does not require introducing the parameter ϵ for the correct definition of such deformed primary fields off criticality. We suppose to describe these problems in other publication.

Acknowledgments

We would like to thank V. Belavin, M. Lashkevich, S. Lukyanov and F. Smirnov for useful discussions. We are grateful to E. Onofri for his kind help in numerical computations. This work was supported by RFBR grant (080100720) and by RBRF-CNRS grant PICS-09-02-91064. Y.P. was supported by Federal Program Scientific and Scientific-Pedagogical Personnel of Innovational Russia under the state contract No. P1339 and by RFBR initiative interdisciplinary project 09-02-12446-ofi-m. The visit of Y.P. to LPTA, University of Montpellier II, was supported by the ENS Landau Exchange Program. Y.P would like to thank the members of the Laboratory and especially A. Neveu, for kind hospitality.

---

[6] Similar problems were independently discussed in the paper [43]. See also Refs. [41, 42].

A Short distance expansion for order and disorder fields

For completeness, we collect here also the results [7] about the following correlation functions

$$G_{+}(x)=\langle\sigma_1(x)\sigma_1^{+}(0)\rangle\,,\quad G_{-}(x)=\langle\mu_1(x)\mu_1^{+}(0)\rangle\,,\tag{87}$$

of order and disorder fields in scaling Z_N Ising model. We considered the leading terms in the short distance expansion

$$G_{\pm}(x)=r^{-4d_1}\left(C_{\pm}^{I}+C_{\pm}^{\varepsilon_1}\langle\varepsilon_1\rangle+C_{\pm}^{\varepsilon_2}\langle\varepsilon_2\rangle+C_{\pm}^{E_1}\langle E_1\rangle+\cdots\right)\,.\tag{88}$$

Up to the first order perturbation theory, the structure functions entering this expression are

$$\begin{aligned}
C_{\pm}^{I}&=1\pm\frac{\lambda\pi r^{2(1-2u)}(\gamma(u)\gamma(3u))^{1/2}\gamma(4u)}{2(1-4u)^2\gamma^3(2u)}\,,\\
C_{\pm}^{\varepsilon_1}&=\mp\frac{r^{4u}(\gamma(u)\gamma(3u))^{1/2}}{2\gamma(2u)}+\frac{\lambda\pi r^2u^2\gamma(4u)\gamma^4(u)}{4\gamma^4(2u)}\,,\\
C_{\pm}^{\varepsilon_2}&=\mp r^{12u}\frac{\lambda\pi r^{2(1-2u)}u^2\gamma^3(u)(\gamma(5u))^{1/2}}{12(1+2u)^2\gamma^2(2u)(\gamma(3u))^{1/2}}\,,\\
C_{\pm}^{E_1}&=-r^{2+4u}\frac{u(1-4u)(\gamma(u)\gamma(3u))^{1/2}}{2(1+2u)(1-2u)\gamma(2u)}\,.
\end{aligned}\tag{89}$$

Note, that the Ising model correlation functions short distance expansions [15, 39] agree with $N\to 2$ expressions of these correlators as well.[7]

B Two particle form factors

Let the contour C goes from infinity above the real axe, then around zero, and then to infinity, below the real axe. Introduce the notation

$$S_2(x)=\exp\frac{1}{2}\int_C\frac{dt}{2\pi it}\frac{\sinh(x-2\pi)t}{\sinh^2(\pi t)}\log(-t)\,.\tag{90}$$

Then the functions $\zeta^{(\dagger)}$, appearing in the equation (33), read

$$\begin{aligned}
\zeta(\beta)&=\frac{i\sinh(\frac{\beta}{2})}{2\sinh(\frac{\beta}{2}+\frac{i\pi}{N})\sinh(\frac{\beta}{2}-\frac{i\pi}{N})}\times\\
&\qquad\qquad\frac{S_2(i\beta+2\pi+\frac{2\pi}{N})S_2(-i\beta+\frac{2\pi}{N})}{S_2^2(2\pi+\frac{2\pi}{N})}\,,\\
\zeta^{\dagger}(\beta)&=\frac{1}{\cosh\frac{\beta}{2}}\frac{S_2^2(2\pi+\frac{2\pi}{N})}{S_2(i\beta+3\pi+\frac{2\pi}{N})S_2(-i\beta+\pi+\frac{2\pi}{N})}\,.
\end{aligned}\tag{91}$$

---

[7] The same phenomena was observed by V. Belavin in the case [40] of the minimal models of the CFT $\mathcal{M}_{2,2n+1}$ perturbed first energy field.

C Integral J_N

In this Appendix we discuss briefly the problem of the calculation of the integral $J_N = \frac{1}{\pi}\int G(z,\overline{z})d^2z$, (78), where

$$G(z,\overline{z}) = \left\langle \Phi^2_{2,0}\,(0)\,\Phi^2_{-2,0}\,(1)\,\epsilon_1(z,\overline{z})\epsilon_1(\infty)\right\rangle_{CFT}. \tag{92}$$

Using a free field representation for parafermionic CFT (see for example [44, 8]), we can express this correlation function in terms of four dimensional integrals. It is convenient to apply the integral transformation derived in Refs. [45] and represent $G(z,\overline{z})$ in the form

$$G(z,\overline{z}) = |z(1-z)|^{-8u}\,(\mathcal{N}_1\; g_1(z,\overline{z}) - \mathcal{N}_2\; g_2(z,\overline{z})). \tag{93}$$

The numerical coefficients $\mathcal{N}_1$ and $\mathcal{N}_2$ are normalization factors. Their exact values are given as (we use the notation $u = \frac{1}{N+2}$)

$$\mathcal{N}_2 = \frac{4(1-2u)^2\gamma(4u)\gamma(5u)}{\pi^2u^2\gamma^3(2u)\gamma(3u)}, \quad \mathcal{N}_1 = -\mathcal{N}_2\frac{5u}{2(1-2u)}\,, \tag{94}$$

The functions $g_1(z,\overline{z})$ and $g_2(z,\overline{z})$ in the equation (93) are defined by the integrals:

$$g_1(z,\overline{z}) = \int d^2t_1d^2t_2\; Q(t_1,t_2,z), \qquad g_2(z,\overline{z}) = \int d^2t_1d^2t_2\; R(t_1,t_2,z), \tag{95}$$

where

$$R(t_1,t_2,z) = |t_1t_2(1-t_1)(1-t_2)(z-t_1)(z-t_2)|^{2u}\,|t_1-t_2|^{4u-4}\;, \tag{96}$$

$$Q(t_1,t_2,z) = R(t_1,t_2,z)\frac{(\overline{t_1}-\overline{t_2})^2}{(\overline{t_1}-\overline{z})(\overline{t_2}-\overline{z})}\,. \tag{97}$$

It follows from the results of paper [46], that the functions $g_1(z,\overline{z})$ and $g_2(z,\overline{z})$ satisfy third order differential equations for each of the variables z and $\overline{z}$. Namely,

$$\begin{array}{c} L_1(z)g_1(z,\overline{z}) = L_1(\overline{z})g_1(z,\overline{z}) = 0\,, \\ L_2(z)g_1(z,\overline{z}) = L_2(\overline{z})g_2(z,\overline{z}) = 0\,. \end{array}$$

Here the differential operators $L_1(z)$ and $L_2(z)$ can be written as polynomials in the parameter u. For $L_1(z)$ we have the definition

$$\begin{aligned} L_1(z) \;\;=\;\; & z(1-z)\partial_z^2 z(1-z)\partial_z + u(8z(1-z)(2z-1)\partial_z^2 + l_1^{(1)}\partial_z) + \\ & + u^2(l_1^{(2)}\partial_z + 8(1-2z)) + u^3 48(1-2z)\,, \end{aligned} \tag{98}$$

with

$$
\begin{aligned}
l_1^{(1)} &= 2(z^2 + (1-z)^2 - 11z(1-z))\,, \\
l_1^{(2)} &= 12z^2 + 12(1-z)^2 - 52z(1-z)\,.
\end{aligned}
$$

The operator $L_2(z)$ is defined in a similar form as following

$$
\begin{aligned}
L_2(z) \quad = \quad & \partial_z z(1-z)\partial_z z(1-z)\partial_z + u(8z(1-z)(2z-1)\partial_z^2 + l_2^{(1)}\partial_z) + \\
& + u^2(l_2^{(2)}\partial_z + 8(1-2z)) + u^3 48(1-2z)\,. \qquad (99)
\end{aligned}
$$

The coefficients $l_2^{(1)}$ and $l_2^{(2)}$ are

$$
\begin{aligned}
l_2^{(1)} &= -8(z^2 + (1-z)^2) + 38z(1-z)\,, \\
l_2^{(2)} &= l_1^{(2)}.
\end{aligned}
$$

Contrary to the case when correlation functions satisfy second order (hypergeometric) differential equation we do not know at present how to calculate in the analytic form the integrals associated with correlation functions satisfying third order equations. However, it is possible to calculate them numerically. The simplest way to do this is to develop a series expansion for the solutions convergent inside and outside a circle $|z| = 1$. We can represent integral J_N in the form $J_N = J_N^{(0)} + J_N^{(\infty)}$, where

$$
J_N^{(0)} = \frac{1}{\pi}\int_{|z|<1} G(z,\overline{z})d^2z, \quad J_N^{(\infty)} = \frac{1}{\pi}\int_{|z|>1} G(z,\overline{z})d^2z. \qquad (100)
$$

Here we consider the integral $J_N^{(0)}$. The calculation of integral $J_N^{(\infty)}$ follows the same steps.

We choose the basis of solutions to the eqs (98,99) y_i, w_i in the form:

$$
y_i(z) = z^{\nu_i(y)}\left(1 + \sum_{j=1} d_j^{(i)}(y)z^j\right),\ w_i(z) = z^{\nu_i(w)}\left(1 + \sum_{j=1} d_j^{(i)}(w)z^j\right). \qquad (101)
$$

It follows from equations (98, 99) that $\nu_1(y) = \nu_1(w) = 0$, $\nu_2(y) = \nu_2(w) + 1 = 1 + 2u$, $\nu_3(y) = \nu_3(w) = 6u$. All the coefficients $d_j^{(i)}(y)$ $d_j^{(i)}(w)$ can be derived from the corresponding differential equations. These coefficients determine the coefficients in the expansion of the "conformal blocks"

$$
Y_i(z) = z^{-4u}(1-z)^{-4u}y_i = z^{-4u+\nu_i(y)}\left(1 + \sum_{j=1} d_j^{(i)}(Y)z^j\right), \qquad (102)
$$

and similar coefficients $d_j^{(i)}(W)$ in the expansion of $z^{-4u}(1-z)^{-4u}w_i$. The value of the integral $J_N^{(0)}$ can be expressed in terms of these coefficients as:

$$\begin{aligned} J_N^{(0)} &= \frac{1}{\pi}\sum_{i=1}^{3}\int_{|z|<1} C_i(Y_i(z)W_i(\overline{z}) - r_iY_i(z)Y_i(\overline{z}))d^2z = \qquad (103) \\ &\sum_{i=1}^{3} C_i\Bigg(\sum_{j=0}^{\infty}\frac{d_j^{(i)}(Y)d^{(i)}_{j+\nu_i(y)-\nu_i(w)}(W) - r_id_j^{(i)}(Y)d_j^{(i)}(Y)}{j+1-4u+\nu_i(y)}\Bigg). \end{aligned}$$

Here the structure constants C_i and numbers r_i can be easily derived from eqs. (93,95) as the coefficients before the corresponding singularities at $z \to 0$ in the correlation function $G(z,\overline{z})$. They are equal:

$$C_1 = \frac{1}{2},\ C_2 = -\frac{u(1-2u)\gamma^2(4u)\gamma^3(u)}{(1+2u)(1-4u)\gamma^4(2u)\gamma(3u)},\ C_3 = \frac{5\gamma(5u)\gamma(u)}{9\gamma^2(3u)} \qquad (104)$$

and $r_1 = 1$, $r_2 = -\frac{2u(1-2u)}{(1+2u)(1-4u)}$, $r_3 = \frac{2}{5}$.

For completeness we give here the expression for $J_N^{(\infty)}$. To write the solutions outside the unit circle it is convenient to introduce the variable $s = 1/z$ and to do the same substitution in the differential operators L_1 and L_2. The operator L_1 is invariant under this substitution and operator L_2 transforms to another operator $\widetilde{L}_2$. The corresponding "conformal blocks", $\widetilde{Y}_i(s) = (1-s)^{-4u}y_i(s)$ and $\widetilde{W}_i(s) = (1-s)^{-4u}\widetilde{w}_i(s)$ will have the following expansion

$$\widetilde{Y}_i(s) = s^{\nu_i(y)}\Bigg(1+\sum_{j=1} d_j^{(i)}(Y)s^j\Bigg),\quad \widetilde{W}_i(s) = s^{\nu_i(\widetilde{w})}\Bigg(1+\sum_{j=1} d_j^{(i)}(\widetilde{W})s^j\Bigg), \qquad (105)$$

where the exponents $\nu_i(y)$ were given above and $\nu_1(\widetilde{w}) = \nu_1(y)$, $\nu_2(\widetilde{w}) = \nu_2(y)$, $\nu_3(\widetilde{w}) = \nu_3(y) + 2$.

The integral $J_N^{(\infty)}$ can be expressed in terms of the coefficients $d_j^{(i)}(Y)$ and $d_j^{(i)}(\widetilde{W})$ as

$$J_N^{(\infty)} = \sum_{i=1}^{3} C_i\Bigg(\sum_{j=0}^{\infty}\frac{q_id_j^{(i)}(Y)d^{(i)}_{j+\nu_i(y)-\nu_i(w)}(\widetilde{W}) - r_id_j^{(i)}(Y)d_j^{(i)}(Y)}{j-1+\nu_i(y)}\Bigg), \qquad (106)$$

where numbers r_i were given above and $q_1 = 3$, $q_2 = -\frac{4u}{(1+2u)}$, $q_3 = \frac{2u^2}{(1+3u)(1+5u)}$.

We note that "dangerous" term with $i = j = 1$ in the sum (106) does not contribute because $d_1^{(1)}(Y) = d_1^{(1)}(\widetilde{W}) = 0$ and the first two terms for $i = 3$ containing $d_{-2}^{(3)}(\widetilde{W})$ and $d_{-1}^{(3)}(\widetilde{W})$ also do not contribute because these coefficients vanish. The infinite sums (103,106) for small u converge rather fast. The terms in these sums decrease as $j^{-\alpha}$ where $\alpha = \min\{4-8u, 3+4u\}$.

The solutions y_i, w_i and $\widetilde{w}_i$ as well as the corresponding differential operators admit the expansions in term of parameter $u = \frac{1}{N+2}$. As an illustration we give here the expansion of functions y_1, w_1 and $\widetilde{w}_1$ up to $O(u^4)$.

$$
\begin{aligned}
y_1 &= 1+4u\log(1-z)+12u^2\log^2(1-z)+\\
&\quad u^3(24\log^3(1-z)-16\log(1-z)PL(2,z)-8PL^{(2)}(2,z))+O(u^4),\\
w_1 &= 1+8u\log(1-z)+u^2(28\log^2(1-z)-8PL(2,z))+\\
&\quad u^3(\frac{176}{3}\log^3(1-z)-48\log(1-z)PL(2,z)-16PL(3,z))+O(u^4),\\
\widetilde{w}_1 &= 1+4u\log(1-z)+\frac{32}{3}u^2\log^2(1-z)+\\
&\quad u^3(\frac{184}{9}\log^3(1-z)-\frac{32}{3}(\log(1-z)PL(2,z)+PL^{(2)}(2,z)))+O(u^4).
\end{aligned}
$$

The functions $PL(2,z)$, $PL(3,z)$, $PL^{(2)}(2,z)$ are $PL(2,z)=-\int_0^z \frac{dt}{t}\log(1-t)$, $PL(3,z)=\int_0^z \frac{dt}{t}PL(2,t)$ and $PL^{(2)}(2,z)=\int_0^z \frac{dt}{t}\log^2(1-t)$.

Using u expansion for the solutions y_i, w_i and $\widetilde{w}_i$ we were able to find explicitly five first terms in the expansion of integral J_N

$$
J_N=\frac{18}{5}u(1+2u+\frac{44}{3}u^2+a_1u^3+a_2u^4)+O(u^6)\,, \tag{107}
$$

where the coefficients a_1 and a_2 have the form

$$
\begin{aligned}
a_1 &= \frac{4}{3}(31\pi^2-2(65+57\zeta(3)))\,,\\
a_2 &= -\frac{8}{3}(234-32\pi^2+\pi^4-67\zeta(3)-92\zeta(5))\,.
\end{aligned}
$$

We can now construct the function $J_N^{(pade)}$, which has the expansion (107), coincides with I_3 given by eq. (80) at $N=3$ and has the asymptotic (82) at $N\to 4$.

$$
J_N^{(pade)}=\frac{4u}{3(1-6u)}+\frac{34u(1-1.032u+14.312u^2-129.08u^3)}{15(1-0.679u+11.953u^2-78.82u^3)}\,. \tag{108}
$$

The function $J_N^{(pade)}$ can be compared with the values $J_N^{(num)}$, which were calculated numerically by Enrico Onofri [47]

N	$J_N^{(pade)}$	$J_N^{(num)}$
3	-1.2109	-1.2109
5	1.6046	1.6046
7	0.67566	0.67522
10	0.4031	0.4033
20	0.184705	0.184721
50	0.0723001	0.0723002
100	0.03603784	0.03603788

(109)

As it is seen from the table, an approximation of the integral J_N for $N\geq 3$ by the function $J_N^{(pade)}$ is done with a very good accuracy. The maximum deviation between data is less then 0.06 per cents.

At the end of this Appendix we note, that it is possible to analyze in a similar spirit another integral, appearing in CFT perturbative calculations

$$I_N = \frac{1}{\pi}\int d^2z\ 2|z(1-z)|^{-8u}\ \mathcal{N}_2\ g_2(z,\overline{z})\,, \tag{110}$$

(see eqs. (93,95))). This expression appears as an integral of correlation function of four thermal fields ϵ_1. It determines the first perturbative correction to the structure constant $C^{\epsilon_1}_{\epsilon_1,\epsilon_1}$

$$I_N = \frac{1}{\pi}\int \langle\epsilon_1(0)\epsilon_1(1)\epsilon_1(z)\epsilon_1(\infty)\rangle_{CFT}\, d^2z. \tag{111}$$

The integral has an series expansion

$$I_N = \frac{24}{5}u(1+2u+8u^2+b_1u^3+b_2u^4)+O(u^6)\,, \tag{112}$$

where the coefficients b_1 and b_2 are given as

$$\begin{aligned} b_1 &= 2(31\pi^2-220-54\zeta(3))\,,\\ b_2 &= 2(196-34\pi^2+2\pi^4-67\zeta(3)-95\zeta(5))\,. \end{aligned}$$

The expression for $I_N^{(pade)}$ can be derived by using the same conditions as for calculation of the $J_N^{(pade)}$, besides the condition that $I_3^{(pade)}$ coincides with $I_3 = 10.0897$. However the value $I_3^{(pade)} = 10.0914$ is very close to I_3. We obtain the following answer

$$I_N^{(pade)} = -\frac{8u}{3(1-6u)}+\frac{8u}{(1-4u)}-\frac{8u(1+13.053u-11u^2+107.2u^3)}{15(1+1.053u-11.585u^2)}\,. \tag{113}$$

For $N \geq 3$ this function approximates I_N calculated numerically [47] with an accuracy which is better then 0.03 per cents

N	$I_N^{(pade)}$	$I_N^{(num)}$
5	-0.2462	-0.2462
7	0.56130	0.56124
10	0.4640	0.4640
20	0.240352	0.240348
50	0.096099	0.096099
100	0.0480157	0.0480157

(114)

References

[1] Al. B. Zamolodchikov. Two point correlation function in scaling Lee-Yang model. *Nucl. Phys.*, **B348**, 619-641 (1991).

[2] A. B. Zamolodchikov. Thermodynamic Bethe ansatz in relativistic models. Scaling three state Potts and Lee-Yang models. *Nucl.Phys.* **B342** 695-720 (1990).

[3] Al. B. Zamolodchikov. Mass scale in the sine-Gordon model and its reductions. *Int.J.Mod.Phys.* **A10**, 1125-1150 (1995).

[4] S. Lukyanov, A. Zamolodchikov. Exact expectation values of local fields in the quantum sine-Gordon model. *Nucl. Phys.* **B493**, 571-587 (1997);
V. A. Fateev, S. L. Lukyanov, A. B. Zamolodchikov, and Al. B. Zamolodchikov. Expectation values of local fields in the Bullough-Dodd model and integrable perturbed Conformal Field Theories. *Nucl. Phys.* **B516**, 652-674 (1998);
V. Fateev, D. Fradkin, S. Lukyanov, A. Zamolodchikov, and Al. Zamolodchikov. Expectation values of descendent fields in the sine-Gordon model. *Nucl. Phys.* **B540**, 587-609 (1999).

[5] M. Karowski, P. Weisz. Exact form factors in (1+1)-dimensional field theoretic models with soliton behavior. *Nucl. Phys.* **B139**, 455476 (1978).

[6] F. A. Smirnov. Form factors in completely integrable models of quantum field theory. Adv. Series in Math. Phys. 14. Singapore: World Scientific (1992).

[7] V. A. Fateev, V. V. Postnikov, Y. P. Pugai. On scaling fields in Z_N Ising models. *JETP Lett.* **83**, 172-178 (2006).

[8] V. A. Fateev, Y. P. Pugai. Expectation values of scaling fields in Z(N) Ising models. *Theor.Math.Phys.* **154**, 473-494 (2008); *Teor.Mat.Fiz.* **154**, 557-583 (2008).

[9] A. B. Zamolodchikov, V. A. Fateev. Parafermionic currents in the two-dimensional conformal quantum field theory and selfdual critical points in Z(N) invariant statistical systems, *Sov. Phys. JETP*, **62**(2), 215225 (1985).

[10] A. B. Zamolodchikov, V. A. Fateev. Disorder fields in two-dimensional conformal quantum field theory and N=2 extended supersymmetry. *Sov. Phys. JETP*, **63**, 913-919 (1986).

[11] V. A. Fateev. Integrable deformations in Z(N) symmetrical models of conformal quantum field theory. *Int. J. Mod. Phys.*, **A6**, 2109-2132 (1991).

[12] R. Koberle, J. A. Swieca. Factorizable Z(N) models. *Phys. Lett.* **B86**, 209210 (1979).

[13] A. B. Zamolodchikov. Integrable field theory from conformal field theory. *Adv. Stud. in Pure Math.* **19**, 641-674 (1989).

[14] J. L. Cardy, G. Mussardo. Form-Factors Of Descendent Operators In Perturbed Conformal Field Theories. *Nucl. Phys.* **B340**, 387-402 (1990);
F. A. Smirnov. Counting the local fields in SG theory. *Nucl.Phys.* **B453**, 807-824 (1995);

A. Koubek. The Space of local operators in perturbed conformal field theories. *Nucl.Phys.* **B435**, 703-734 (1995);
M. Jimbo, T. Miwa, Y Takeyama. Counting minimal form-factors of the restricted sine-Gordon model. math-ph/0303059.

[15] T. T. Wu, B M. McCoy, C. A. Tracy, E. Barouch. Spin spin correlation functions for the two-dimensional Ising model: Exact theory in the scaling region. *Phys.Rev.* **B13**, 316-374 (1976).

[16] F.C. Alcaraz, R. Koberle. Duality and the phases of Z(N) spin systems. *J.Phys.* **A13**, L153 (1980).

[17] L. P. Kadanoff and H. Ceva. Determination of an operator algebra for the two-dimensional ising model. *Phys. Rev.* **B11**, 3918-3939 (1971).

[18] E. Fradkin, L. P. Kadanoff. Disorder variables and para-fermions in two-dimensional statistical mechanics. *Nucl. Phys.* **B170 [FS1]**, 1 (1980).

[19] A. A. Belavin, A. M. Polyakov, A. B. Zamolodchikov. Infinite conformal symmetry in two-dimensional Quantum Field Theory. *Nucl. Phys.* **B241**, 333-380 (1984).

[20] F.C. Alcaraz. Discrete mass spectrum of Z(N) spin systems perturbed by a thermal field. *J.Phys.* **A23**, L1105-L1108 (1990).

[21] V. A. Fateev. The exact relations between the coupling constants and the masses of particles for the integrable perturbed conformal field theories. *Phys. Lett.* **B324**, 45-51 (1994).

[22] O. Babelon, D. Bernard, F. A. Smirnov. Null vectors in integrable field theory. *Commun.Math.Phys.* **186**, 601-648 (1997).

[23] M. Jimbo, H. Konno, S. Odake, Y. Pugai, and J. Shiraishi. Free field construction for the ABF models in Regime II. *J. Statist. Phys.* **102**, 883921 (2001).

[24] M. Jimbo, T. Miwa, Algebraic analysis of lattice models. Conference Board of the Mathematical Sciences. AMS.

[25] S. Lukyanov, Y. Pugai. Bosonization of ZF algebras: Direction toward deformed Virasoro algebra. *J. Exp. Theor. Phys.* **82**, 1021-1045 (1996); Multipoint local height probabilities in the integrable RSOS model. *Nucl. Phys.* **B473**, 631-658 (1996).

[26] Y. Pugai. Vacuum expectation values from fusion of vertex operators. *JETP Lett.* **79**, 457 (2004); *Pisma Zh.Eksp.Teor.Fiz.* **79**, 569-574 (2004).

[27] A. B. Zamolodchikov. Infinite additional symmetries in two-dimensional conformal Quantum Field Theory. *Theor. Math. Phys.* **65**, 1205-1213 (1985);

V. A. Fateev, A. B. Zamolodchikov. The models of two-dimensional conformal field theory having Z(3) symmetry. *Nucl.Phys.* **B280**, 644-660 (1987).
V. Fateev, S. Lukyanov. The models of two-dimensional conformal Quantum Field Theory with Z(n) symmetry. *Int. J. Mod. Phys.* **A3**, 507 (1988).

[28] V. A. Fateev and A. B. Zamolodchikov. Selfdual solutions of the star triangle relations in Z(N) models. *Phys. Lett.*, **92A**, 37-39, (1982).

[29] F. A. Smirnov. Quantum groups and generalized statistics in integrable models. *Comm. Math. Phys.*, **132**, 415-439, (1990);
A. N. Kirillov, F. A. Smirnov. preprint ITF-88-73R, Kiev (1988).

[30] H. Babujian, A. Foerster, M. Karowski. Exact form factors in integrable quantum field theories: the scaling Z(N)-Ising model. *Nucl.Phys.* **B736**, pp. 169-198 (2006);
H. Babujian, M. Karowski, Exact form factors for the scaling ZN-Ising and the affine A_{N-1}-Toda quantum field theories. *Phys.Lett.* **B575**, 144-150 (2003).

[31] M. Caselle, G. Delfino, P. Grinza, O. Jahn and N. Magnoli. Potts correlators and the static three-quark potential. *J. Stat. Mech.* **0603**, P008 (2006).

[32] G. E. Andrews, R. J. Baxter, P. J. Forrester. Eight-vertex SOS model and generalized Rogers-Ramanujan-type identities *J. Stat. Phys.* **35**, 193 (1984).

[33] S. Lukyanov. Free field representation for massive integrable models. *Comm. Math. Phys.* **167**, 183-226 (1995); Form-factors of exponential fields in the sine-Gordon model. *Mod. Phys. Lett.* **A12**, 2543-2550 (1997); Form-factors of exponential fields in the affine A(N)(1)-1 Toda model. *Phys. Lett.*, **B408**, 192-200 (1997).

[34] S. L. Lukyanov, A. B. Zamolodchikov. Form-factors of soliton creating operators in the sine-Gordon model. *Nucl.Phys.* **B607**, 437-455 (2001).

[35] V. A. Fateev. Normalization factors, reflection amplitudes and integrable systems. e-Print: hep-th/0103014; Normalization factors in conformal field theory and their applications. *Mod. Phys. Lett.* **A15**, 259-270 (2000).

[36] P. Fonseca, A. Zamolodchikov. Ward identities and integrable differential equations in the Ising field theory. e-Print: hep-th/0309228.

[37] V.S. Dotsenko, V.A. Fateev. Conformal algebra and multipoint correlation functions in two-dimensional statistical models. *Nucl. Phys.* **B240**[**FS12**], 312(1984); Four point correlation functions and the operator algebra in the two-dimensional conformal invariant theories with the central charge $c < 1$. *Nucl. Phys.* **B251**[**FS13**], 691 (1985).

[38] P. Baseilhac, M. Stanishkov. Expectation values of descendent fields in the Bullough-Dodd model and related perturbed conformal field theories. *Nucl.Phys.* **B612**, 373-390 (2001).

[39] A. B. Zamolodchikov Expectation value of composite field T anti-T in two-dimensional quantum field theory. e-Print: hep-th/0401146.

[40] A. A. Belavin, V. A. Belavin, A. V. Litvinov, Y. P. Pugai, Al. B. Zamolodchikov. On correlation functions in the perturbed minimal models M(2 , 2n+1). *Nucl.Phys.* **B676**, 587-614 (2004).

[41] V.A. Belavin, O.V. Miroshnichenko. Correlation functions of descendants in the scaling Lee–Yang model. *JETP Lett.* **82**,679-684 (2005).

[42] G. Delfino, G. Niccoli. Form-factors of descendant operators in the massive Lee-Yang model. *J.Stat.Mech.* **0504**, P004 (2005).

[43] B. Feigin, M. Lashkevich. Form factors of descendant operators: Free field construction and reflection relations. *J.Phys.A* **42** P. 304014 (2009). arXiv:0812.4776.

[44] D. Nemeshansky, *Phys. Lett.* **B224**, 121 (1989)

[45] V.A. Fateev, A.V. Litvinov. Multipoint correlation functions in Liouville field theory and minimal Lioville gravity. *Theor. Math. Phys.* **154**, 454-472 (2008); arXiv hep-th 0707.1664; V.A. Fateev, A.V. Litvinov. Coulomb integrals in Liouville field theory and Lioville gravity. *JETP Lett.* **84**, 531-536 (2007).

[46] V. A. Fateev, A.V. Litvinov, A.Neveu, E.Onofri. Differential equation for four-point correlation function and elliptic four-point conformal bloks. *J. Phys.* **A42**, 304011, 2009; arXiv hep-th:0902.1331.

[47] E. Onofri, *Unpublished.*